## QUATERNARY GEOLOGY

## LIST OF MAP UNITS



hydrography, geographic names and geographic places. Streets and street names are from Tele Atlas® copyrighted data. Base map information is neither current nor complete. CONTOUR DATA - Derived from Connecticut's 2000 statewide LiDAR, (Light Detection And Ranging), dataset by the University of Connecticut, College of Agriculture and Natural Resources, Department of

Natural Resources and the Environment. These data are a Beta product intended for research and demonstration purposes. NOTE: Contour line data is known to be incorrect in some areas due to anomalies in the underlying elevation data used to generate those specific contour lines. Areas where contour lines are too straight or angular, do not naturally curve where expected, or don't exist where they probably should are good indications of erroneous data.

> Map is not colorfast Protect from light and moisture



Map created by CT DEP December 2010

STATE OF CONNECTICUT DEPARTMENT OF ENVIRONMENTAL PROTECTION 79 Elm Street Hartford, CT 06106-5127

IAP UNITS					
L <b>TWATER DEPOSITS - late Wisconsinan</b> ntiated Meltwater Deposits	Explanation of Map Symbols				Palmer
of Major Ice-Dammed Lakes of Major Sediment-Dammed Lakes	Ice Margin Position Inferred Ice Margin Position				
of Related Series of Major Ice-Dammed Ponds of Related Series of Major Sediment-Dammed Ponds	<ul><li>Section Strict Strict Strict</li><li>Glacial Striction or Groove</li></ul>				
of Proximal Meltwater Streams	<ul><li>Drumlin Axis and Center</li><li>Meltwater Channel</li></ul>				
LAID DEPOSITS - late Wisconsinan, Illinoian Deposits	→ Glacial Lake Spillway → Inferred Glacial Spillway				
Deposits ine Deposits	<ul> <li>Location of Lower Till</li> <li>Two-Till Outcrop</li> </ul>				
ours	Deltaic Bedding Locality     Weathered Bedrock Outcrop				
terval erval	<ul> <li>Radiocarbon-Dated Locality</li> </ul>				
IATION					
Glacial Meltwater Deposits (sorted and stratified deltaic, rideposits); and Postglacial Deposits (flood-plain alluvium and terrace, talus, dune, tidal-marsh, beach, channel fill, marine	ver bottom, lake bottom, and inland dune swamp deposits, but also including stream- delta deposits, and artificial fill) that were				
emplaced in comparable topographic and depositional settings are categorized and color coded in the Legend Description. Re axes, ice-margin positions, scarps, drainage divides, gla	s, and therefore share similar characteristics, lated Map Elements include eskers, drumlin cial lake spillways, meltwater channels,				
striations/grooves, dated sample locations, glaciofluvial and la map units and various types of exposures.	ke-bottom facies as overlays on glacial lake				
<b>Glacial Ice-Laid Deposits</b> (nonsorted and generally nonstra were derived directly from the ice and consist of nonsorted, ge ranging from clay to large boulders. The matrix of most tills i	tified thin till, thick till, and end moraine) enerally nonstratified mixtures of grain-sizes is predominantly sand and silt, and boulders				
can be sparse to abundant. Some tills contain lenses of sorted laminated fine-grained sediment. The lack of sorting and stra makes them poorly drained, difficult to dig in or plow, mediou	sand and gravel and occasionally masses of atification typical of ice-laid deposits often cre sources of groundwater and unsuited for				
septic systems. Till blankets the bedrock surface in variable th meltwater deposits. End moraine deposits (primarily ablat Connecticut. Ice-laid deposits are inferred to be of Wiscons	icknesses and commonly underlies stratified ion till) occur principally in southeastern sinan age except where exposures of older				
(probably Illinoian) till are shown. Drumlins are inferred to be till.	e composed of older till mantled by younger				
Glacial Meltwater Deposits (sorted and stratified deltaic, r deposits) were laid down in glacial streams, lakes and ponds Connecticut as the last ice sheet systematically (Koteff and P	which occupied the valleys and lowlands of ressl, 1981) melted away to the north. They				
boulders, and owing to their water-related depositional origination for development. Because water is a better sorting a	ins they have many characteristics that are gent than ice, glacial meltwater deposits are				
sources of construction aggregate, and are relatively easy to ex Stratified meltwater deposits include both fine and coarse gr	cavate and build highways and buildings on. rained deposits such as silt, clay, sand, and				
The mapping presented here and on the Quaternary Geology Basin is based on recognizing single bodies of sediment or ass	Map of Connecticut and Long Island Sound				
can be identified as mappable units known as morphoseque sedimentary facies are associated with fluvial, deltaic and lake are emplaced in high-energy settings at or near the ice front F	ences (Koteff and Pessl, 1981). Different e-bottom settings. Coarse proximal deposits				
the glacier (distally) and grain size decreased along the morphosequences are coarse grained at their collapsed, ice-com 1). A detailed discussion of the complexities and significant	e path of meltwater flow. As a result, tact heads and become finer distally (Figure ce of morphosequences is contained in the	len			
pamphlet that accompanies the Quaternary Geology Map of Co	onnecticut and Long Island Sound Basin.	Hamp			
Lake-B	Bottom				
and forms, grading from ice-contact forms (eskers, kames) to non-ice rgin of a glacier, graded to a specific base level. Grain-size decreas	e-contact forms es from coarse				
et slopes to silt and clay in lake-bottom deposits (after Stone and othe	prs, 2005).				
sequences occupying higher, narrower portions of the valle opposite is true. The ice itself was the impoundment, and the o	y (Figure 2). In north-draining systems the dest morphosequences were emplaced in the				
spillways were opened and the valleys widened. In this case, t the lowest, widest portions of the valley (Figure 2).	he youngest depositional sequences occupied				
	South				
Narrow Basin					
ing Basin					
			Tray		
Wide Basin		for (			Sawmill Brook
					State Still Brook
			Scantic River		Porta
ing Dasin at-dammed basins (Bottom). The mechanism of impoundment an ve to the direction of ice retreat. These relationships are reflected	d the chronological I in the organization	a		Crow Hill Brook	
				Sta	A Start
resources. Talus, a result of rockfall at the base of steep bed dune deposits, that developed as winds swept across newly ex-	rock (primarily trap rock) cliffs, and inland cposed glacial lake beds, provide ecological				The second of th
elements of coastal and poorly drained inland settings. De composed of sands, gravels and silts that have been reword	eposits of floodplain alluvium are largely ked from glacial deposits and mixed with			C Hill C A Start	32 S I A
have historically been attractive for agricultural uses and commerce.	development related to water-dependant		Spencer.Rd Pond		
OURCES				~ ) / / / / / / / /	
RELATED INFORMATION		5755			Parking
I HIS map IS Intended to be printed at its original dimensions, (3 scale (1 inch = 2,000 feet).	x + x + 30 in), in order to maintain the 1:24,000	5 2	Chesthut Hill		Dam Middle Lave Steen
data of Connecticut Quaternary Geology and Surficial Materia the Connecticut Department of Environmental Protection, in content and the data were digitized from the 1:24,000 cools are in the	als combined into one dataset, published by operation with the U.S. Geological Survey.		Ellis Brook		Pond Dam
Materials Map of Connecticut, (Stone, J.R., Schafer, J.P., Lon Geological Survey Special Map, 2 sheets, scale 1:125,000, ma Geologic Map of Connecticut and Long Island Sound Basin	don, E.H. and Thompson, W.B., 1992, U.S. ap and pamphlet, 71 p.) and the Quaternary (Stone, J.R., Schafer, I.P. London, F.H.				Stafford Springs
DiGiacomo-Cohen, M.L., Lewis, R.L., and Thompson, W.B Investigation Map 2784, 2 sheets, scale 1:125,000).	., 2005, U.S. Geological Survey Scientific		MAP LOCATION		
OTHER GEOLOGIC MAPS - This map is also available for in intended to be used with other bedrock, surficial, and quaterr reports published by the Connecticut Geological and Natural	ndividual towns of Connecticut. This map is nary (glacial) geology quadrangle maps and History Survey, USGS, and others. Those			10.5	0
maps are reports are also available from CT DEP. MAPS AND DIGITAL DATA - Go to the CT ECO website fo	r this map and a variety of others. Go to the			1000 0 1000	2000 3000 4000 5000 6000 7
y CT DEP				10.5 SCALE 1:24,000 (1 inch	= 2,000 feet) when map is printed at original s
r 2010 🔀			State Plane Coordinate System of 1983, Zone 3526		

Lambert Conformal Conic Projectio North American Datum of 1983



MONSON, CONNECTICUT CT DEP Quadrangle 10