







## LIST OF MAP UNITS

Explanation of Map Symbols	Elevation Contours	Weathered Bedrock Outcrop
 Area of glaciofluvial deposits grading to glacial lake	 100 Ft. Interval	 Weathered Bedrock Outcrop
 Area of lake-bottom sediments	50 Ft. Interval	 Radiocarbon-Dated Locality
Drainage Divide - Boundary between major geologic basins.		
 Drainage Divide - Boundary within major geologic basin dividing it into north-draining and south-draining regions		

Quaternary Geology is 1:24,000-scale data that illustrates the geographic features formed in Connecticut during the Quaternary Period, which spans from 2.588 ± 0.005 millions years ago to the present and includes the Pleistocene (glacial) and Holocene (postglacial) Epochs. The Quaternary Period has been a time of development of many depositional features, including glacial drift, alluvial fans, and eskers. In the last Pleistocene, continental ice sheets swept across Connecticut from the north. Their effects are of pervasive importance to present-day occupants of the land.

The Connecticut Geological Survey information was initially compiled at 1/24,000 scale (1 inch = 2,000 feet) then recompiled for a statewide 1/25,000-scale map. The Quaternary Geological Map of Connecticut and Long Island Sound Basin. A companion map of the Quaternary Geology of the Connecticut Coastal Plain. The maps show the subsurface textures, grain-size distribution of the materials. The quaternary geology and surficial material features portrayed on these two maps are very closely related; each contributes to the interpretation of the other.

Particular attention has been paid to understanding the distribution and characteristics of stratified meltwater deposits because they have historically influenced development patterns and groundwater availability throughout the state. Within the meltwater category, six classes of deposits have been recognized based on the conditions that created them after emplacement. Four of these are based on the previous depositional environment, or the glacier itself, impounded the lake or pond where emplacement occurred (see the meltwater deposit discussion below). Meltwater stream deposits are defined by the distance from the glacier front to the deposit, and are subdivided when they were emplaced, and a separate meltwater map unit is reserved for deposits of undetermined provenance (uncorrelated).

Glacial Ice-Laid Deposits (nonsorted and generally nonstratified thin till, thick till, and end moraine); Glacial Meltwater Deposits (sorted and stratified deltaic, river bottom, accompanies the Quaternary Geology Map of Connecticut and Long Island Sound Basin.

**Figure 1:** A morphosequence is a body of meltwater deposits composed of a continuum of land forms, grading from ice-contact forms (eskers, kames) to non-ice-contact forms (flat valley terrace, delta plains), that were deposited simultaneously at and beyond the margin of a glacier, grading to a specific base level. Grain-size decreases from coarse gravel at ice-contact heads, through sand and gravel and sand beneath delta plains and foreset slopes to silt and clay in lake-bottom deposits (after Stone and others, 2005).

North Wide Basin Narrow Basin South

**Figure 2:** Scenario for morphosequence development in ice-dammed (Top) and sediment-dammed basins (Bottom). The mechanism of impoundment and the chronological and topographic positions of the deposits are related to the orientation of the basins relative to the direction of ice retreat. These relationships are reflected in the organization and color coding of the List of Map Units (after Stone and others, 2005).

Postglacial deposits provide locally important ecological, agricultural, commercial, prone nature, low, flat, fertile floodplains have historically been attractive for agricultural uses and development related to water-dependant commerce.

**QUATERNARY GEOLOGY DATA** – Quaternary Geology shown on this map are from the Quaternary Geology Poly, Point Feature, and Line Feature dataset intended to be used at 1:24,000 scale. Based on Connecticut Quaternary Geology digital spatial data published in 2005 by the U.S. Geological Survey, in cooperation with the Connecticut Department of Environmental Protection. These data were digitized from the 1:24,000 scale compilation sheets prepared for the Geological and Quaternary Geology Map of Connecticut, (Stone, J.R., Schaller, J.P., London, E.H., Digiomo-Cohen, M. L., Lewis R.S. and Thompson, W.B., 2005, U.S. Geological Survey special map, 2 sheets, scale 1:125,000).

**CONTOUR DATA** - Derived from Connecticut's 2000 statewide LiDAR (Light Detection and Ranging) data by the University of Connecticut, College of Agriculture and Natural Resources, Department of Natural Resources and the Environment. These data are a beta product intended for research and demonstration purposes. NOTE: Contour line data is known to be incorrect in some areas due to anomalies in the underlying elevation data used to generate those specific contour lines. Areas where contour lines are too straight or angular, do not naturally curve where expected, or don't exist where they probably should are good indicators of erroneous data.

**OTHER GEOLOGIC MAPS** - Digital map data is also available for individual USGS topographic quadrangles of Connecticut. This map is intended to be used with other bedrock, surficial, and quaternary (glacial) geology town maps and reports published by the Connecticut Geological and Natural History Survey, USGS, and others. Those maps are reports are also available from CT DEP.

**MAPS AND DIGITAL DATA** - Go to the CT ECHO website for this map and a variety of others. Go to the CT DEP website for the digital spatial data shown on this map.

MAP LOCATION


0 0.25 0.5 1 1.5 2 Miles

0 1.125 2.250 4.500 6.750 8.000 Feet

0 0.25 0.5 1 1.5 2 Kilometers


SCALE 1:24,000 (1 inch = 2,000 feet) when map is printed at original size (48 x 36 in)

State Plane Coordinate System of 1983, Zone 3526  
Lambert Conformal Conic Projection  
North American Datum of 1983


 STATE OF CONNECTICUT  
 DEPARTMENT OF  
 ENVIRONMENTAL PROTECTION  
 79 Elm Street  
 Hartford, CT 06106-5127

Map created by CT DEP  
 December 2010

Map is not colorfast  
 Protect from light and moisture


 USGS  
*science for a changing world*




















---

[illegible]

favorable for development. Because water is a better sorting agent than ice, glacial meltwater deposits are commonly better sorted, more permeable, and better aquifers than ice-laid deposits. They can be good sources of construction aggregate, and are relatively easy to excavate and build highways and buildings on. Stratified meltwater deposits include both fine and coarse grained deposits such as silt, clay, sand, and gravel.

The mapping presented here and on the Quaternary Geology Map of Connecticut and Long Island Sound Basin is based on recognizing single bodies of sediment or assemblages of glacial sedimentary facies that can be identified as mappable units known as morphosequences (Kotěff and Pessl, 1981). Different sedimentary facies are associated with different glacial and/or ice-sheet systems, and these sedimentary facies are

associated with moraine, deltaic and lake-bottom settings. Coarse proximal deposits are displaced in high-energy settings at or near the ice front. Energy levels dropped off with distance from the glacier (distally) and grain size decreased along the path of meltwater flow. As a result, morphosequences are coarse grained at their collapsed, ice-contact heads and become finer distally (Figure 1). A detailed discussion of the complexities and significance of morphosequences is contained in the pamphlet that

accompanies the Quaternary Geology Map of Connecticut and Long Island Sound basin.

forms, grading from ice-contact forms (eskers, kames) to non-ice-contact forms (flat valley bedded to a specific base level. Grain-size decreases from coarse gravel at ice-contact heads, to fine sand and silt in the middle of the valley, to clay in the tail deposits (after Stone and others, 2005).

progressed up valley, with the youngest depositional sequences occupying higher, narrower portions of the valley (Figure 2). In north-draining systems the opposite is true. The ice itself was the impoundment, and the oldest morphosequences were emplaced in the higher, narrower portions of the basin. As the ice front retreated northward, a succession of lower bedrock spillways were opened and the valleys widened. In this case, the youngest depositional sequences occupied the lowest, widest portions of the valley (Figure 2).

South  
Narrow Basin

basins (Bottom). The mechanism of impoundment and the chronological and topographic retreat. These relationships are reflected in the organization and color coding of the List of

and recreational resources. Talus, a result of rockfall at the base of steep bedrock (primarily trap rock) cliffs, and inland dune deposits, that developed as winds swept across newly exposed glacial lake beds, provide ecological niches that are atypical for Connecticut. Beach, dune, marsh and swamp deposits are key ecological elements of coastal and poorly drained inland settings. Deposits of floodplain alluvium are largely composed of sands, gravels and silts that have been reworked from glacial deposits and mixed with organic matter which increases their fertility. Despite their flood-

prone nature, low, flat, fertile floodplains have historically been attractive for agricultural uses and development related to water-dependant commerce.

## SOURCES

**RELATED INFORMATION**

This map is intended to be printed at its original dimensions, (48 x 36 in), in order to maintain the 1:24,000 scale (1 inch = 2,000 feet).

**QUATERNARY GEOLOGY AND SURFICIAL MATERIALS DATA - 1:24,000-scale digital spatial data of Connecticut Quaternary Geology and Surficial Materials combined into one dataset, published by the Connecticut Department of Environmental Protection, in cooperation with the U.S. Geological Survey. These data were digitized from the 1:24,000-scale compilation sheets prepared for both the**

Sedimental Materials Map of Connecticut (Stone, J.R., Schärer, J.P., London, E.H. and Thompson, W.B., 1992, U.S. Geological Survey Special Map, 2 sheets, scale 1:125,000, map and pamphlet, 71 p.) and the Quaternary Geologic Map of Connecticut and Long Island Sound Basin (Stone, J.R., Schärer, J.P., London, E.H., DiGiacomo-Cohen, M.L., Lewis, R.L., and Thompson, W.B., 2005, U.S. Geological Survey Scientific Investigation Map 2784, 2 sheets, scale 1:125,000).

**OTHER GEOLOGIC MAPS** - This map is also available for individual USGS topographic quadrangles of Connecticut. This map is intended to be used with other bedrock, surficial, and quaternary (glacial) geology town maps and reports published by the Connecticut Geological and Natural History Survey, USGS, and others. Those maps are reports are also available from CT DEP.

**MAPS AND DIGITAL DATA** - Go to the CT ECO website for this map and a variety of others. Go to the CT DEP website for the digital spatial data shown on this map.

MAP LOCATION

2 miles  
6 in

State Plane Coordinate System of 1983, Zone 3526  
Lambert Conformal Conic Projection  
North American Datum of 1983

### Discussion

USGS  
science for a changing world

---

