

LIST OF MAP UNITS


EXPLANATION

Figure 1: A morphosequence is a body of meltwater deposits composed of a continuum of (terrace, delta plains), that were deposited simultaneously at and beyond the margin of a glacier through sand and gravel and sand beneath delta plains and foreset slopes to silt and clay in lake.

Figure 2: Scenario for morphosequence development in ice-dammed (Top) and sediment-dammed (Bottom) positions of the deposits are related to the orientation of the basins relative to the direction

Postglacial deposits provide locally important ecological, agricultural, commercial,

SCALE 1:24,000 (1 inch = 2,000 feet) when map is printed at original size



STATE OF CONNECTICUT
DEPARTMENT OF
ENVIRONMENTAL PROTECTION
79 Elm Street
Hartford, CT 06106-5127

The mapping presented here and on the Quaternary Geology Map of Connecticut and Long Island Sound Basin is based on recognizing single bodies of sediment or assemblages of glacial sedimentary facies that can be identified as mappable units known as morphosquences (Kottief and Pessl, 1981). Different sedimentary facies are associated with fluvial, deltaic and lake-bottom settings. Coarse proximal deposits are associated with energy-saturated, high velocity, high discharge, high sedimentation rate, and with distance from the glacier (distally) and grain size decreased along the path of meltwater flow. As a result, morphosquences are coarse grained at their collapsed, ice-contact heads and become finer distally (Figure 1). A detailed discussion of the complexities and significance of morphosquences is contained in the pamphlet that accompanies the Quaternary Geology Map of Connecticut and Long Island Sound Basin.

South
Lanka Battaram

forms, grading from ice-contact forms (eskers, kames) to non-ice-contact forms (flat valley graded to a specific base level. Grain-size decreases from coarse gravel at ice-contact heads, bottom deposits (after Stone and others, 2005).

progressed up valley, with the youngest depositional sequences occupying higher, narrower portions of the valley (Figure 2). In north-draining systems the opposite is true. The ice itself was the impoundment, and the oldest morphosequences were emplaced in the higher, narrower portions of the basin. As the ice front retreated northward, a succession of lower bedrock spillways were opened and the valleys widened. In this case, the youngest depositional sequences occupied the lowest, widest portions of the valley (Figure 2).

South



ing Basin

Geological cross-section of the Rio de Janeiro area. The diagram shows a series of rock units dipping to the right. From left to right, the units are: a thick layer of yellowish-brown material (likely sedimentary or volcanic), a red layer (likely igneous or metamorphic), and a grey layer (likely igneous or metamorphic). Below these units is a wavy line representing the base of the sequence. The text 'Rio de Janeiro' is partially visible on the left side of the diagram.

d basins (Bottom). The mechanism of impoundment and the chronological and topographic retreat. These relationships are reflected in the organization and color coding of the List of

and recreational resources. Talus, a result of rockfall at the base of steep bedrock (primarily trap rock) cliffs, and inland dune deposits, that developed as winds swept across newly exposed glacial lake beds, provide ecological niches that are atypical for Connecticut. Beach, dune, marsh and swamp deposits are key ecological elements of coastal and poorly drained inland settings. Deposits of floodplain alluvium are largely composed of sands, gravels and silts that have been reworked from glacial deposits and mixed with organic matter which increases their fertility. Despite their flood-

agricultural uses and development related to water-dependant commerce.

RELATED INFORMATION
This map is intended to be printed at its original dimensions, (48 x 36 in), in order to maintain the 1:24,000 scale (1 inch = 2,000 feet).

QUATERNARY GEOLOGY AND SURFICIAL MATERIALS DATA - 1:250,000-scale digital spatial data of Connecticut Quaternary Geology and Surficial Materials combined into one dataset, published by the Connecticut Department of Environmental Protection, in cooperation with the U.S. Geological Survey. These data were digitized from the 1:250,000-scale compilation sheets prepared for both the Surficial Materials Map of Connecticut, (Stone, J.R., Schafer, J.P., London, E.H., and DiGiacomio-Cohen, M.L., Lewis, R.L., and Thompson, W.B., 2005, U.S. Geological Survey Scientific Investigation Map 2784, 2 sheets, scale 1:250,000).

OTHER GEOLOGIC MAPS - This map is also available for individual USGS topographic quadrangles of Connecticut. This map is intended to be used with other bedrock, surficial, and quaternary (glacial) geology town maps and reports published by the Connecticut Geological and Natural History Survey, USGS, and others. Those maps and reports are also available from CT DEP.

MAPS AND DIGITAL DATA - Go to the CT ECO website for this map and a variety of others. Go to the CT DEP website for the digital spatial data shown on this map.

MAP LOCATION

0 100 Miles

State Plane Coordinate System of 1983, Zone 3626
Lambert Conformal Conic Projection
North American Datum of 1983

