## QUATERNARY GEOLOGY LYME, CONNECTICUT LIST OF MAP UNITS **Explanation of Map Symbols** POSTGLACIAL DEPOSITS - late Holocene, late Wisconsinan GLACIAL MELTWATER DEPOSITS - late Wisconsinan Artificial Fill Ice Margin Position Undifferentiated Meltwater Deposits **— — —** Inferred Ice Margin Position Coastal Beach and Dune Deposits Deposits of Major Ice-Dammed Lakes Deposits of Major Sediment-Dammed Lakes Tidal Marsh Deposits → Glacial Striation or Groove Deposits of Related Series of Major Ice-Dammed Ponds Floodplain Alluvium Drumlin Axis and Center Deposits of Related Series of Major Sediment-Dammed Ponds Swamp Deposits Meltwater Channel Deposits of Proximal Meltwater Streams ── Glacial Lake Spillway EARLY POSTGLACIAL DEPOSITS - early Holocene, late Wisconsinan Deposits of Distal Meltwater Streams Inferred Glacial Spillway GLACIAL ICE-LAID DEPOSITS - late Wisconsinan, Illinoian Stream Terrace Deposits ♦ Location of Lower Till Thin Till Deposits Inland Dune Deposits ♦ Two-Till Outcrop Thick Till Deposits ▲ Deltaic Bedding Locality End Moraine Deposits × Weathered Bedrock Outcrop **Explanation of Map Symbols Elevation Contours** Radiocarbon-Dated Locality Area of glaciofluvial deposits grading to glacial lake 100 Ft. Interval Area of lake-bottom sediments 50 Ft. Interval Drainage Divide -- Boundary between major geologic basins. ---- Drainage Divide -- Boundary within major geologic basin EASTHADDAM dividing it into north-draining and south-draining regions **EXPLANATION** Quaternary Geology is 1:24,000-scale data that illustrates the geologic features formed lake bottom, and inland dune deposits); and Postglacial Deposits (flood-plain alluvium in Connecticut during the Quaternary Period, which spans from 2.588 ± 0.005 million and swamp deposits, but also including stream-terrace, talus, dune, tidal-marsh, beach, years ago to the present and includes the Pleistocene (glacial) and Holocene channel fill, marine delta deposits, and artificial fill) that were emplaced in comparable (postglacial) Epochs. The Quaternary Period has been a time of development of many topographic and depositional settings, and therefore share similar characteristics, are details of the Connecticut landscape and all surficial deposits. At least twice in the last categorized and color coded in the Legend Description. Related Map Elements include eskers, drumlin axes, ice-margin positions, scarps, drainage divides, glacial lake Pleistocene, continental ice sheets swept across Connecticut from the north. Their spillways, meltwater channels, striations/grooves, dated sample locations, glaciofluvial effects are of pervasive importance to present-day occupants of the land. and lake-bottom facies as overlays on glacial lake map units and various types of The Quaternary Geology information illustrates the geologic history and the distribution of depositional environments during the emplacement of unconsolidated glacial and postglacial surficial deposits and the landforms resulting from those events Glacial Ice-Laid Deposits (nonsorted and generally nonstratified thin till, thick till, and in Connecticut. These deposits range from a few feet to several hundred feet in end moraine) were derived directly from the ice and consist of nonsorted, generally thickness, overlie the bedrock surface and underlie the organic soil layer of nonstratified mixtures of grain-sizes ranging from clay to large boulders. The matrix of Connecticut. Quaternary Geology is mapped without regard for any organic soil layer most tills is predominantly sand and silt, and boulders can be sparse to abundant. Some tills contain lenses of sorted sand and gravel and occasionally masses of laminated finegrained sediment. The lack of sorting and stratification typical of ice-laid deposits often The Connecticut Quaternary Geology information was initially compiled at 1:24,000 makes them poorly drained, difficult to dig in or plow, mediocre sources of scale (1 inch = 2,000 feet) then recompiled for a statewide 1:125,000-scale map, groundwater and unsuited for septic systems. Till blankets the bedrock surface in Quaternary Geology Map of Connecticut and Long Island Sound Basin. A companion variable thicknesses and commonly underlies stratified meltwater deposits. End map, the Surficial Materials Map of Connecticut, emphasizes the surface and moraine deposits (primarily ablation till) occur principally in southeastern Connecticut. MONTVILLE subsurface texture (grain-size distribution) of these materials. The quaternary geology Ice-laid deposits are inferred to be of Wisconsinan age except where exposures of older (probably Illinoian) till are shown. Drumlins are inferred to be composed of older till and surficial material features portrayed on these two maps are very closely related; each contributes to the interpretation of the other. mantled by younger till. Most of Connecticut's surficial material is glacially derived, and can be divided into Glacial Meltwater Deposits (sorted and stratified deltaic, river bottom, lake bottom, two broad depositional categories: Glacial Ice-Laid Deposits (nonsorted and generally and inland dune deposits) were laid down in glacial streams, lakes and ponds which nonstratified thin till, thick till, and end moraine) which are generally exposed in the occupied the valleys and lowlands of Connecticut as the last ice sheet systematically uplands, and are the most widespread surficial deposit in Connecticut; and Glacial (Koteff and Pessl, 1981) melted away to the north. They are often composed of layers dune deposits) which are most commonly concentrated in valleys and lowlands. owing to their water-related depositional origins they have many characteristics that are favorable for development. Because water is a better sorting agent than ice, glacial Particular attention has been paid to understanding the distribution and characteristics meltwater deposits are commonly better sorted, more permeable, and better aquifers of stratified meltwater deposits because they have historically influenced development than ice-laid deposits. They can be good sources of construction aggregate, and are patterns and groundwater availability throughout the state. Within the meltwater relatively easy to excavate and build highways and buildings on. Stratified meltwater category, six classes of deposits have been recognized based on the conditions that deposits include both fine and coarse grained deposits such as silt, clay, sand, and prevailed during their emplacement. Four of the seven indicate whether previously gravel. deposited sediment, or the glacier itself, impounded the lake or pond where emplacement occurred (see the meltwater deposit discussion below). Meltwater stream The mapping presented here and on the Quaternary Geology Map of Connecticut and deposits are differentiated based on their distance (proximal or distal) from the ice sheet Long Island Sound Basin is based on recognizing single bodies of sediment or when they were emplaced, and a separate meltwater map unit is reserved for deposits of assemblages of glacial sedimentary facies that can be identified as mappable units undetermined provenance (uncorrelated). known as morphosequences (Koteff and Pessl, 1981). Different sedimentary facies are associated with fluvial, deltaic and lake-bottom settings. Coarse proximal deposits are **Postglacial Deposits** were emplaced by various processes after the melt back of the last emplaced in high-energy settings at or near the ice front. Energy levels dropped off ice sheet. Some of these deposits were emplaced early in post-glacial time and have with distance from the glacier (distally) and grain size decreased along the path of been grouped together as Early Postglacial Deposits. Later deposits, resulting from meltwater flow. As a result, morphosequences are coarse grained at their collapsed, iceprocesses that are still active (or are manmade), have been grouped together as contact heads and become finer distally (Figure 1). A detailed discussion of the complexities and significance of morphosequences is contained in the pamphlet that accompanies the Quaternary Geology Map of Connecticut and Long Island Sound **Glacial Ice-Laid Deposits** (nonsorted and generally nonstratified thin till, thick till, and Basin. end moraine); Glacial Meltwater Deposits (sorted and stratified deltaic, river bottom, Ice-Contact (collapsed) Figure 1: A morphosequence is a body of meltwater deposits composed of a continuum of land forms, grading from ice-contact forms (eskers, kames) to non-ice-contact forms (flat valley terrace, delta plains), that were deposited simultaneously at and beyond the margin of a glacier, graded to a specific base level. Grain-size decreases from coarse gravel at ice-contact heads, through sand and gravel and sand beneath delta plains and foreset slopes to silt and clay in lake-bottom deposits (after Stone and others, 2005). Deposition of the morphosequences that progressively filled bedrock valleys and progressed up valley, with the youngest depositional sequences occupying higher, lowlands as the last glacier melted northward required the presence of impounded narrower portions of the valley (Figure 2). In north-draining systems the opposite is lakes and ponds. The nature of the impoundments and the resulting distribution of the true. The ice itself was the impoundment, and the oldest morphosequences were meltwater deposits on the landscape were controlled by the topography of the area emplaced in the higher, narrower portions of the basin. As the ice front retreated being deglaciated. Where a northward succession of ice positions was established in northward, a succession of lower bedrock spillways were opened and the valleys south-draining basins, previously deposited sediment formed the dams, and the oldest widened. In this case, the youngest depositional sequences occupied the lowest, widest morphosequences occupied the lowest, widest parts of the valley. Deposition then North-Draining Basin South-Draining Basin Figure 2: Scenario for morphosequence development in ice-dammed (Top) and sediment-dammed basins (Bottom). The mechanism of impoundment and the chronological and topographic positions of the deposits are related to the orientation of the basins relative to the direction of ice retreat. These relationships are reflected in the organization and color coding of the List of Postglacial Deposits (flood-plain alluvium and swamp deposits, but also including and recreational resources. Talus, a result of rockfall at the base of steep bedrock stream-terrace, talus, dune, tidal-marsh, beach, channel fill, marine delta deposits, and (primarily trap rock) cliffs, and inland dune deposits, that developed as winds swept artificial fill) are less widely distributed and are typically thinner than the glacial across newly exposed glacial lake beds, provide ecological niches that are atypical for Bushy Hill Pond deposits that they overlie. The oldest postglacial deposits occur in Long Island Sound Connecticut. Beach, dune, marsh and swamp deposits are key ecological elements of and in southeastern Connecticut because these areas were deglaciated first. Many of coastal and poorly drained inland settings. Deposits of floodplain alluvium are largely ESSE composed of sands, gravels and silts that have been reworked from glacial deposits the depositional processes that were initiated as postglacial conditions began to and mixed with organic matter which increases their fertility. Despite their floodprevail are still operative today. prone nature, low, flat, fertile floodplains have historically been attractive for Postglacial deposits provide locally important ecological, agricultural, commercial, agricultural uses and development related to water-dependant commerce. DATA SOURCES QUATERNARY GEOLOGY DATA – Quaternary Geology shown on this map are from the Quaternary Geology Poly, Point Feature, and Line Feature dataset intended This map is intended to be printed at its original dimensions, (48 x 36 in), in order to to be used at 1:24,000 scale. Based on Connecticut Quaternary Geology digital spatial maintain the 1:24,000 scale (1 inch = 2,000 feet). data published in 2005 by the U.S. Geologic Survey, in cooperation with the Connecticut Department of Environmental Protection. These data were digitized QUATERNARY GEOLOGY AND SURFICIAL MATERIALS DATA - 1:24,000from the 1:24,000-scale compilation sheets prepared for the statewide Quaternary scale digital spatial data of Connecticut Quaternary Geology and Surficial Materials Geology Map of Connecticut, (Stone, J.R., Schafer, J.P., London, E.H., DiGiacomocombined into one dataset, published by the Connecticut Department of Cohen, M. L., Lewis R.S, and Thompson, W.B., 2005, U.S. Geological Survey Environmental Protection, in cooperation with the U.S. Geological Survey. These special map, 2 sheets, scale 1:125,000). data were digitized from the 1:24,000-scale compilation sheets prepared for both the Surficial Materials Map of Connecticut, (Stone, J.R., Schafer, J.P., London, E.H. and Thompson, W.B., 1992, U.S. Geological Survey Special Map, 2 sheets, scale BASE MAP DATA - Based on data originally from 1:24,000-scale USGS 7.5 minute 1:125,000, map and pamphlet, 71 p.) and the Quaternary Geologic Map of topographic quadrangle maps published between 1969 and 1992. It includes political boundaries, railroads, airports, hydrography, geographic names and geographic Connecticut and Long Island Sound Basin, (Stone, J.R., Schafer, J.P., London, E.H., places. Streets and street names are from Tele Atlas® copyrighted data. Base map DiGiacomo-Cohen, M.L., Lewis, R.L., and Thompson, W.B., 2005, U.S. Geological Survey Scientific Investigation Map 2784, 2 sheets, scale 1:125,000). information is neither current nor complete. CONTOUR DATA - Derived from Connecticut's 2000 statewide LiDAR, (Light OTHER GEOLOGIC MAPS - This map is also available for individual USGS topographic quadrangles of Connecticut. This map is intended to be used with other Detection And Ranging), dataset by the University of Connecticut, College of Agriculture and Natural Resources, Department of Natural Resources and the bedrock, surficial, and quaternary (glacial) geology town maps and reports published by the Connecticut Geological and Natural History Survey, USGS, and others. Those Environment. These data are a Beta product intended for research and demonstration purposes. NOTE: Contour line data is known to be incorrect in some areas due to maps are reports are also available from CT DEP. anomalies in the underlying elevation data used to generate those specific contour MAPS AND DIGITAL DATA - Go to the CT ECO website for this map and a lines. Areas where contour lines are too straight or angular, do not naturally curve where expected, or don't exist where they probably should are good indications of variety of others. Go to the CT DEP website for the digital spatial data shown on this erroneous data. OLD SAYBROOK WESTBROOK SCALE 1:24,000 (1 inch = 2,000 feet) when map is printed at original size (48 x 36 in) State Plane Coordinate System of 1983, Zone 3526 Lambert Conformal Conic Projection North American Datum of 1983 TATE OF CONNECTICUT Map created by CT DEP DEPARTMENT OF December 2010 ENVIRONMENTAL PROTECTION Map is not colorfast

Hartford, CT 06106-5127

Protect from light and moisture