

LIST OF MAP UNITS

Quaternary Geology is 1:24,000-scale data that illustrates the glacial history formed in Connecticut during the Quaternary Period, which spans from 2.588 ± 0.005 million years ago to the present and includes both the Pleistocene (time of glacial) and Holocene (postglacial) epochs. The Quaternary Period is the time of development of many deposits of the Connecticut landscape, and surficial deposits exist in the last 100,000 years. Pleistocene, continental ice sheets swept across Connecticut from the north. Their effects are pervasive importance to present-day occupants of the land.

The Quaternary Geology information illustrates the glacial history and the distribution of depositional environments during the emplacement of unconsolidated glacial and postglacial surficial deposits and the landforms resulting from those events in Connecticut. These deposits range from a few feet to several hundred feet in thickness, and they underlie the glacial and postglacial ice layer of the Connecticut. Quaternary Geology is mapped without regard for any organic soil layer that may overlie the deposit.

Quaternary Geology is 1:24,000-scale data that illustrates the glacial history formed in Connecticut during the Quaternary Period, which spans from 2.588 ± 0.005 million years ago to the present and includes both the Pleistocene (time of glacial) and Holocene (postglacial) epochs. The Quaternary Period is the time of development of many deposits of the Connecticut landscape, and surficial deposits exist in the last 100,000 years. Pleistocene, continental ice sheets swept across Connecticut from the north. Their effects are pervasive importance to present-day occupants of the land.

The Quaternary Geology information illustrates the glacial history and the distribution of depositional environments during the emplacement of unconsolidated glacial and postglacial surficial deposits and the landforms resulting from those events in Connecticut. These deposits range from a few feet to several hundred feet in thickness, and they underlie the glacial and postglacial ice layer of the Connecticut. Quaternary Geology is mapped without regard for any organic soil layer that may overlie the deposit.


Most of Connecticut's surficial material is glacially derived, and can be divided into two broad depositional categories: Glacial Ice-Laid Deposits (nonsorted and generally nonstratified thin till, thick till, and end moraine) which are generally exposed in the uplands, and are the most widespread surficial deposit in Connecticut; and Glacial Meltwater Deposits (sorted and stratified deltaic, river bottom, lake bottom, and inland dune deposits) which are most commonly concentrated in valleys and lowlands.

Postglacial Deposits were emplaced by various processes after the melt back of the last ice sheet. Some of these deposits were emplaced early in post-glacial time and have been grouped together as Early Postglacial Deposits. Later deposits, resulting from processes that are still active (or are manmade), have been grouped together as Recent Deposits.

Postglacial deposits provide locally important ecological, agricultural, commercial,

QUATERNARY GEOLOGY DATA – Quaternary Geology shown on this map are from the Quaternary Geology Poly, Point Feature, and Line Feature dataset intended to be used at 1:24,000 scale. Based on Connecticut Quaternary Geology digital spatial data published in 2005 by the U.S. Geologic Survey, in cooperation with the Connecticut Department of Environmental Protection. These data were digitized from the 1:24,000-scale compilation sheets prepared for the statewide Quaternary Geology Map of Connecticut, (Stone, J.R., Schafer, J.P., London, E.H., DiGiacomo, Cohen, M. L., Lewis R.S. and Thompson, W.B., 2005, U.S. Geological Survey special map, 2 sheets, scale 1:125,000).

CONTOUR DATA - Derived from Connecticut's 2000 statewide LiDAR, (Light Detection And Ranging), dataset by the University of Connecticut, College of Agriculture and Natural Resources, Department of Natural Resources and the Environment. These data are a Beta product intended for research and demonstration purposes. NOTE: Contour line data is known to be incorrect in some areas due to anomalies in the underlying elevation data used to generate those specific contour lines. Areas where contour lines are too straight or angular, do not naturally curve where expected, or don't exist where they probably should are good indications of erroneous data.


 STATE OF CONNECTICUT
 DEPARTMENT OF
 ENVIRONMENTAL PROTECTION
 79 Elm Street
 Hartford, CT 06106-5127

Map created by CT DEH
December 2010
Map is not colorfast
Protect from light and moisture

WATER DEPOSITS - late Wisconsin

- ▲ at Melhatch Deposits
- Major Ice-Dammed Lakes
- Major Settlement-Dammed Lakes
- Related Series of Major Ice-Dammed Ponds
- Related Series of Major Settlement-Dammed Ponds
- Provincial Melhatch Streams
- Provincial Melhatch Streams

ICE DEPOSITS - late Wisconsin, Illinoian

- deposits
- deposits
- Ice Deposits

Glacial Ice-Laid Deposits (nonsorted and generally nonstratified thin till, thick till, and moraine) were derived directly from the ice and consist of nonsorted, generally nonstratified mixtures of grain-size ranging from clay to large boulders. The matrix of most tills is predominantly sand and silt, and boulders can be sparse to abundant. Some tills contain lenses of silt and gravel and occasionally masses of laminated fine-grained sediment. The lack of sorting and stratification typical of ice-laid deposits often results in poorly drained, difficult to dig in or plow, midwestern sources of ground-water and unsuited for septic systems. Till blankets the bedrock surface in variable thicknesses and commonly underlies stratified meltwater deposits. Ice-moraine deposits (primarily ablation till) occur principally in southeastern Connecticut. Ice-laid deposits are inferred to be of Wisconsinan age except where exposures of older (probably Illinoian) till are shown. Drumlins are inferred to be composed of older till remobilized by younger till.

The mapping presented here and on the Quaternary Geology Map of Connecticut and the Long Island Sound Basin is based on recognizing single bodies of sediment or assemblages of glacial sedimentary facies that can be identified as mappable units known as morphosquences (Kottief and Peissl, 1981). Different sedimentary facies are

Basin.

SOURCES

RELATED INFORMATION
This map is intended to be printed at its original dimensions, (48 x 36 in), in order to maintain the 1:24,000 scale (1 inch = 2,000 feet).

OTHER GEOLOGIC MAPS - This map is also available for individual USGS topographic quadrangles of Connecticut. This map is intended to be used with other bedrock, surficial, and quaternary (glacial) geology town maps and reports published by the Connecticut Geological and Natural History Survey, USGS, and others. Those maps are reports are also available from CT DEP.

MAPS AND DIGITAL DATA - Go to the CT ECO website for this map and a variety of others. Go to the CT DEP website for the digital spatial data shown on this map.

by CT DEP
er 2010

colorfast
ht and moisture

 **USGS**
science for a changing world

