

LIST OF MAP UNITS

Quaternary Geology is 1:24,000-scale local map that illustrates the geologic features formed in Connecticut during the Quaternary Period, which spans from 2,588 + 0.005 million years ago to the present. The map shows the distribution of Quaternary glacial and postglacial (postglacial) deposits. The Quaternary Period has been a time of development of many details of the Connecticut landscape and all surficial deposits. At least twice in the last 100,000 years, the Connecticut River valley and the western part of the state north. These effects are of pervasive importance to present-day occupants of the land.

The Quaternary Geology information illustrates the geologic history and the distribution of depositional environments during the emplacement of unconsolidated glacial and postglacial surficial deposits and the landforms resulting from those events. The surficial deposits thickness range from a few feet to several hundred feet in thickness. The surficial deposits range from the glacial till to the organic soil layer. Connecticut. Quaternary Geology is mapped without regard for any organic soil layer that may overlay the deposit.

The Connecticut Quaternary Geology information was initially compiled at 1:24,000 scale (1 inch = 2,000 feet) then recompiled for a statewide 1:125,000-scale map. The Connecticut Map of Quaternary and Long Island Sound organic soil layer map, the Surficial Materials Map of Connecticut, emphasizes the surface and subsurface texture (grain-size distribution) of the surficial materials. The Quaternary geology and surficial materials maps are very closely related. The two maps are very closely related, each contributes to the interpretation of the other.

Quaternary Geology is 1:24,000-scale local map that illustrates the geologic features formed in Connecticut during the Quaternary Period, which spans from 2,588 + 0.005 million years ago to the present. The map shows the distribution of Quaternary glacial and postglacial (postglacial) deposits. The Quaternary Period has been a time of development of many details of the Connecticut landscape and all surficial deposits. At least twice in the last 100,000 years, the Connecticut River valley and the western part of the state north. These effects are of pervasive importance to present-day occupants of the land.

The Quaternary Geology information illustrates the geologic history and the distribution of depositional environments during the emplacement of unconsolidated glacial and postglacial surficial deposits and the landforms resulting from those events. The surficial deposits thickness range from a few feet to several hundred feet in thickness. The surficial deposits range from the glacial till to the organic soil layer. Connecticut. Quaternary Geology is mapped without regard for any organic soil layer that may overlay the deposit.

The Connecticut Quaternary Geology information was initially compiled at 1:24,000 scale (1 inch = 2,000 feet) then recompiled for a statewide 1:125,000-scale map. The Connecticut Map of Quaternary and Long Island Sound organic soil layer map, the Surficial Materials Map of Connecticut, emphasizes the surface and subsurface texture (grain-size distribution) of the surficial materials. The Quaternary geology and surficial materials maps are very closely related, and both maps are very closely related, both contribute to the interpretation of the other.

Most of Connecticut's surficial material is glacially derived, and can be divided into two broad depositional categories: Glacial Ice-Laid Deposits (nonsorted and generally nonstratified thin till, thick till, and end moraine) which are generally exposed in the uplands, and are the most widespread surficial deposit in Connecticut; and Glacial Melting Deposits (sorted and stratified, river and lake bottom deposits, and inland dune deposits) which are most commonly concentrated in valleys and lowlands.

Particular attention has been paid to understanding the distribution and characteristics of stratified meltwater deposits because they have historically influenced the location of patents and groundwater availability throughout the state. Within the meltwater category, six classes of deposits have been recognized based on the conditions that prevailed during their emplacement. Four of the seven indicate whether previously deposited sediment, or the glacier itself, impounded the lake or pond where the deposits were emplaced. The remaining two classes are based on whether the deposits are differentiated based on their distance (proximal or distal) from the ice sheet when they were emplaced, and a separate meltwater map unit is reserved for deposits of undetermined provenance (unsorted).

Glacial Ice-Laid Deposits (nonsorted and generally nonstratified thin till, thick till, and end moraine); **Glacial Meltwater Deposits** (sorted and stratified deltaic, river bottom,


Figure 1: A morphosequence is a body of meltwater deposits composed of a continuum of terrace, delta plains), that were deposited simultaneously at and beyond the margin of a glacier through sand and gravel and sand beneath delta plains and foreset slopes to silt and clay in lake

Postglacial Deposits (flood-plain alluvium and swamp deposits, but also including stream-terrace, talus, dune, tidal-marsh, beach, channel fill, marine delta deposits, and artificial fill) are less widely distributed and are typically thinner than the glacial deposits that they overlie. The oldest postglacial deposits occur in Long Island Sound and in southeastern Connecticut because these areas were deglaciated first. Many of the depositional processes that were initiated as postglacial conditions began to prevail are still operative today.

DATA

QUATERNARY GEOLOGY DATA – Quaternary Geology shown on this map are from the Quaternary Geology Poly, Point Feature, and Line Feature dataset intended to be used at 1:24,000 scale. Based on Connecticut Quaternary Geology digital spatial data published in 2005 by the U.S. Geological Survey, in cooperation with the Connecticut Department of Transportation. The data were digitized from the 1:24,000-scale compilation sheets prepared for the statewide Quaternary Geology Map of Connecticut, (Stone, J.R., Schafer, J.P., London, E.H., DiGiacomo, Cohen, M., Lewis R.S. and Thompson, W.B., 2005, U.S. Geological Survey special map 1, 2, series scale 1:125,000).

BASE MAP DATA - Based on data originally from 1:24,000-scale USGS 7.5 minute topographic quadrangle maps published between 1969 and 1992. It includes 7.5 minute contour intervals, roads, water features, hydrologic features, geographic names and geographic place names. Streets and street names are from Tele Atlas. Copyrighted data. Base map information is neither current nor from Tele Atlas.



STATE OF CONNECTICUT
DEPARTMENT OF
ENVIRONMENTAL PROTECTION
79 Elm Street
Hartford, CT 06106-5127

Glacial Laminates Deposits (nonstratified and generally nonconsisted thin, till, thick, and fine) were derived directly from the ice and consisted of nonstratified generally nonstratified mixtures of grain-sizes ranging from clay to large boulders. The matrix of the tills is predominantly sand and silt, and clowders can be sparse to abundant. Some of the tills are composed of well-sorted, rounded to subrounded clasts of quartz, granite, and gneiss. The laminates are composed of fine-grained sand, silt, and clay, and are generally well-sorted. The lack of sorting and stratification typical of ice-laden deposits often makes them poorly drained, difficult to dig or plow, meddlesome sources of material for agricultural purposes, and generally undesirable for construction. The variable thicknesses and commonly undulating stratified meddlesome deposits. End moraines are composed of till, and are generally poorly sorted, and are composed of variable deposits are inferred to be of Wisconsin origin. Some exposures of older (probably Illinoian) till are shown. Drumlins are inferred to be composed of older till probably by Younger till.

accompanies the Quaternary Geology Map of Connecticut and Long Island Sound Basin.

progressed up valley, with the youngest depositional sequences occupying higher, narrower portions of the valley (Figure 2). In north-draining systems the opposite is true. The ice itself was the impoundment, and the oldest morphosequences were emplaced in the higher, narrower portions of the basin. As the ice front retreated northward, a succession of lower bedrock spillways were opened and the valleys widened. In this case, the youngest depositional sequences occupied the lowest, widest portions of the valley (Figure 2).

and recreational resources. Talus, a result of rockfall at the base of steep bedrock (primarily trap rock) cliffs, and inland dune deposits, that developed as winds swept across newly exposed glacial lake beds, provide ecological niches that are atypical for Connecticut. Beach, dune, marsh and swamp deposits are key ecological elements of coastal and poorly drained inland settings. Deposits of floodplain alluvium are largely composed of sands, gravels and silts that have been reworked from glacial deposits and mixed with organic matter which increases their fertility. Despite their flood-

SOURCES

RELATED INFORMATION

This map is intended to be printed at its original dimensions, (48 x 36 in), in order to maintain the 1:24,000 scale (1 inch = 2,000 feet).

QUATERNARY GEOLOGY AND SURFICIAL MATERIALS data - 1:24,000-scale digital spatial data of Connecticut Quaternary Geology and Surficial Materials combined into one dataset, published by the Connecticut Department of Environmental Protection, in cooperation with the U.S. Geological Survey. These data were derived from a 1:24,000-scale compilation sheets prepared for both the Surficial Materials Map of Connecticut (Stone, J.R., Schafer, J.P., London, E.H. and Thompson, W.B., 1992, U.S. Geological Survey Special Map, 2 sheets, scale 1:125,000, map and pamphlet, 71 p.) and the Quaternary Geologic Map of Connecticut (Stone, J.R., Schafer, J.P., London, E.H., Thompson, W.B., and DeGiacomo-Cohen, M.L., Lewis, R.L., and Thompson, W.B., 2005, U.S. Geological Survey Scientific Investigation Map 2784, 2 sheets, scale 1:125,000).

MAPS AND DIGITAL DATA - Go to the CT ECO website for this map and a variety of others. Go to the CT DEP website for the digital spatial data shown on this map.

State Plane Coordinate System of 1983, Zone 3526
Lambert Conformal Conic Projection
North American Datum of 1983