

# QUATERNARY GEOLOGY NORWICH, CONNECTICUT

## LIST OF MAP UNITS

<b>POSTGLACIAL DEPOSITS - late Holocene, late Wisconsinan</b>	<b>GLACIAL MELT-WATER DEPOSITS - late Wisconsinan</b>	<b>Explanation of Map Symbols</b>
Artificial Fill	Undifferentiated Meltwater Deposits	Ice Margin Position
Coastal Beach and Dune Deposits	Deposits of Major Ice-Dammed Lakes	Inferred Ice Margin Position
Tidal Marsh Deposits	Deposits of Major Sediment-Dammed Lakes	Esker
Floodplain Alluvium	Deposits of Related Series of Major Ice-Dammed Ponds	Glacial Striation or Groove
Swamp Deposits	Deposits of Related Series of Major Sediment-Dammed Ponds	Drumlin Axis and Center
Talus	Deposits of Proximal Meltwater Streams	Meltwater Channel
<b>EARLY POSTGLACIAL DEPOSITS - early Holocene, late Wisconsinan</b>	Deposits of Distal Meltwater Streams	Glacial Lake Spillway
Stream Terrace Deposits	<b>GLACIAL ICE-LAID DEPOSITS - late Wisconsinan, Illinoian</b>	Inferred Glacial Spillway
Inland Dune Deposits	Thin Till Deposits	Location of Lower Till
	Thick Till Deposits	Two-Till Outcrop
	End Moraine Deposits	Delicate Bedding Locality
		Weathered Bedrock Outcrop
		Radiocarbon-Dated Locality

## EXPLANATION

Quaternary Geology is a 1:24,000-scale data that illustrates the geologic features formed in Connecticut during the Quaternary Period, which spans from 2,588 ± 0.005 million years ago to the present and includes the Pleistocene (glacial) and Holocene (postglacial) Epochs. The Quaternary Period has been a time of development of many details of the Connecticut landscape and all surficial deposits. At least twice in the last Pleistocene, continental ice sheets swept across Connecticut from the north. Their effects are of pervasive importance to present-day occupants of the land.

The Quaternary Geology information illustrates the geologic history and the distribution of depositional environments during the emplacement of unconsolidated glacial and postglacial surficial deposits and the landforms resulting from these events in Connecticut. These deposits range from a few feet to several hundred feet in thickness, overlie the bedrock surface and underlie the organic soil layer of Connecticut. Quaternary Geology is mapped without regard for any organic soil layer that may occur in the deposit.

The Connecticut Quaternary Geology information was initially compiled at a 1:24,000 scale (1 inch = 2,000 feet) then recomputed for a statewide 1:125,000-scale map, Quaternary Geology Map of Connecticut and Long Island Sound Basin. A companion map, the Surficial Materials Map of Connecticut, emphasizes the surface and subsurface texture (grain-size distribution) of these materials. The quaternary geology and surficial material features portrayed on these two maps are very closely related, each contributing to the interpretation of the other.

Most of Connecticut's surficial material is glacially derived, and can be divided into two broad depositional categories: Glacial Ice-Laid Deposits (nonstratified till, thick till, and end moraine) which are generally exposed in the uplands, and are the most widespread surficial deposit in Connecticut; and Glacial Meltwater Deposits (sorted and stratified deltaic, river bottom, lake bottom, and inland dune deposits) which are most commonly concentrated in valleys and lowlands.

Particular attention has been paid to understanding the distribution and characteristics of stratified meltwater deposits because they have historically influenced drainage patterns and groundwater availability throughout the state. Within the meltwater category, six classes of deposits have been recognized based on the conditions that prevail during their emplacement. Four of the seven indicate whether previously deposited sediment, or the glacier itself, impounded the lake or pond where emplacement occurred (see the meltwater deposit discussion below). Meltwater stream deposits are differentiated based on their distance (proximal or distal) from the ice sheet when they were emplaced, and a separate meltwater map unit is reserved for deposits of undetermined provenance (uncorrelated).

Postglacial Deposits were emplaced by various processes after the melt back of the last ice sheet. Some of these deposits were emplaced early in post-glacial time and have been grouped together as Early Postglacial Deposits. Later deposits, resulting from processes that are still active (or are manmade), have been grouped together as Postglacial Deposits.

Glacial Ice-Laid Deposits (nonstratified and generally nonstratified till, thick till, and end moraine), Glacial Meltwater Deposits (sorted and stratified deltaic, river bottom, lake bottom, and inland dune deposits), and Postglacial Deposits (flood-plain alluvium and swamp deposits, but also including stream-terrace, talus, dune, tidal-marsh, beach, channel fill, marine delta deposits, and artificial fill) that were emplaced in comparable topographic and depositional settings, and therefore share similar characteristics, are categorized and color coded in the Legend Description. Related Map Elements include eskers, drumlin axes, ice-margin positions, scarp, drainage divides, glacial lake spillways, meltwater channels, striations/grooves, dated sample locations, glaciofluvial and lake-bottom facies as overlays on glacial lake map units and various types of exposures.




Figure 1: A morphosequence is a body of meltwater deposits composed of a continuum of land forms, grading from ice-contact forms (eskers, kames) to non-ice-contact forms (flat valley terrace, delta plains), that were deposited simultaneously and beyond the margin of a glacier, graded to a specific base level. Grain-size decreases from coarse gravel at ice-contact, roads, through sand and gravel and sand beneath delta plains and forest slopes to silt and clay in lake-bottom deposits (after Stone and others, 2005).




Figure 2: Scenario for morphosequence development in ice-dammed (Top) and sediment-dammed basins (Bottom). The inclusion of impoundment and the chronological and topographic positions of the deposits are related to the orientation of the basin relative to the direction of ice retreat. These relationships are reflected in the organization and color coding of the List of Map Units (after Stone and others, 2005).

Postglacial Deposits (flood-plain alluvium and swamp deposits, but also including stream-terrace, talus, dune, tidal-marsh, beach, channel fill, marine delta deposits, and artificial fill) are less widely distributed and are typically thinner than the glacial deposits that they overlie. The oldest postglacial deposits occur in Long Island Sound and in southeastern Connecticut because these areas were deglaciated first. Many of the depositional processes that were initiated as postglacial conditions began to prevail are still operative today.

Postglacial deposits provide locally important ecological, agricultural, commercial, and recreational resources. Talus, a result of rockfall at the base of steep bedrock (primarily trap rock) cliffs, and inland dune deposits, that developed as winds swept across newly exposed glacial lake beds, provide ecological niches that are attractive to Connecticut. Beach, dune, marsh and swamp deposits are key ecological elements of coastal and poorly drained inland settings. Deposits of flood-plain alluvium are largely composed of sands, gravels and silts that have been reworked from glacial deposits and mixed with organic matter which increases their fertility. Despite their flood-prone nature, low, flat, fertile floodplains have historically been attractive for agricultural uses and development related to water-dependent commerce.

## DATA SOURCES

**QUATERNARY GEOLOGY DATA** - Quaternary Geology shown on this map are from the Quaternary Geology Poly, Point Feature, and Line Feature dataset intended to be used at 1:24,000 scale. Based on Connecticut Quaternary Geology digital spatial data published in 2005 by the U.S. Geological Survey, in cooperation with the Connecticut Department of Environmental Protection. These data were digitized from the 1:24,000-scale compilation sheets prepared for the statewide Quaternary Geology Map of Connecticut, (Stone, J.R., Schuler, J.P., London, E.H., DiGiacomo-Cohen, M.L., Lewis, R.S., and Thompson, W.B., 2005, U.S. Geological Survey special map, 2 sheets, scale 1:125,000).

**BASE MAP DATA** - Based on data originally from 1:24,000-scale USGS 7.5 minute topographic quadrangle maps published between 1969 and 1992. It includes political boundaries, railroads, airports, hydrographic names and geographic features. Streets and street names are from Tele Atlas' copyrighted data. Base map information is neither current nor complete.

**CONTOUR DATA** - Derived from Connecticut's 2000 statewide LIDAR (Light Detection And Ranging) dataset by the University of Connecticut, College of Agriculture and Natural Resources, Department of Natural Resources and the Environment. These data are a beta product intended for research and demonstration purposes. NOTE: Contour lines do not necessarily represent the actual elevation data used to generate these specific contour lines. Areas where contour lines are not straight or angular, do not necessarily curve where expected, or don't exist where they probably should are good indications of erroneous data.

MAP LOCATION: State Plane Coordinate System of 1983, Zone 3628 Lambert Conformal Conic Projection North American Datum of 1983

SCALE 1:24,000 (1 inch = 2,000 feet) when map is printed at original size (48 x 36 in)

Map created by CT DEP December 2010

Map is not colorfast. Project from light and moisture

STATE OF CONNECTICUT DEPARTMENT OF ENVIRONMENTAL PROTECTION 79 Elm Street Hartford, CT 06106-5127

USGS science for a changing world



Map is not colorfast. Project from light and moisture

STATE OF CONNECTICUT DEPARTMENT OF ENVIRONMENTAL PROTECTION 79 Elm Street Hartford, CT 06106-5127

USGS science for a changing world