

LIST OF MAP UNITS

[illegible]

Quantary Geology is 1:24,000-scale data that illustrates the geologic features formed in Connecticut during the Quaternary Period, which spans from 2,588 ± 0.005 million years ago to the present. The Quaternary Period is subdivided into the Pleistocene (postglacial) Epochs. The Quaternary Period has been a time of development of many physical features of the Connecticut landscape and all surficial deposits. At least twice in the last Pleistocene Epoch, the Connecticut River valley was covered by ice from the north. These effects are of pervasive importance to present-day occupants of the land.

The Quaternary Geology information illustrates the geologic history and the

distribution of depositional environments during the emplacement of unconsolidated glacial and postglacial surficial deposits and the landforms resulting from those events in Connecticut. These deposits range from a few feet to several hundred feet in thickness, overlie the bedrock surface and underlie the organic soil layer of Connecticut. Quaternary Geology is mapped without regard for any organic soil layer that may overly the deposit.

The Connecticut Geology information was initially compiled at 1:24,000 scale (1 inch = 2,000 feet) then recompiled for a statewide 1:125,000-scale map. The Map of Connecticut and Long Island Sound Basin. A companion map, *Geological Map of Connecticut and Long Island Sound Basin*, shows the subsurface tectonic and stratigraphic relationships, and the geologic and subsurface textural (grain-size distribution) of these materials. The geology, geologic and subsurface material features portrayed on these two maps are very closely related; each contributes to the interpretation of the other.

Most of Connecticut's surficial material is glacially derived, and can be divided into two broad depositional categories: Glacial Ice-Land Deposits (notsorted and generally nonstratified thin till, thick till, and end moraine) which are generally exposed in the uplands, and are the most widespread surficial deposit in Connecticut; and Glacial Meltwater Deposits (sorted and stratified deltaic, river bottom, lake bottom, and inland dune deposits) were laid down in glacial streams, lakes and ponds which occupied the valleys and lowlands of Connecticut as the last ice sheet systematically (Kottfe and Pessl, 1981) melted away to the north. They are often composed of layers of well-to-poorly sorted sands, gravels, silts and clays with few to no boulders, and owing to their water-related depositional origins they have many characteristics that are

Particular attention has been paid to understanding the distribution and characteristics of stratified meltwater deposits because they have historically influenced development patterns and groundwater availability throughout the state. Within the meltwater category, six classes of deposits have been recognized based on the conditions that formed them: (1) ice-contact sediment, (2) ice-rafted sediment, (3) ice-marginal deposited sediment, or the glacier itself, impounded the lake or pond where emplacement occurred (see the meltwater deposit discussion below). Meltwater stream deposits are further divided into (1) ice-contact (proximal) and (2) distal (where they were emplaced, and a separate meltwater map unit is reserved for deposits of undetermined provenance (uncorrelated).

Postglacial Deposits were emplaced by various processes after the melt back of the last ice sheet. Some of these deposits were emplaced early in post-glacial time and have been grouped together as Early Postglacial Deposits. Later deposits, resulting from processes that are still active (or are manmade), have been grouped together as Postglacial Deposits.

Glacial Ice-Laid Deposits (nonsorted and generally nonstratified thin till, thick till, and end moraine); **Glacial Meltwater Deposits** (sorted and stratified deltaic, river bottom,

Deposition of the morphosquences that progressively filled bedrock valleys and lowlands as the last glacier melted northward required the presence of impounded lakes and ponds. The nature of the impoundments and the resulting distribution of the meltwater deposits on the landscape were controlled by the topography of the area being deglaciated. Where ice stagnated in the north, the largest, most extensive north-draining basins, previously deposited sediment formed the dams, and the oldest morphosquences occupied the lowest, widest parts of the valley. Deposition then

Figure 2: Scenario for morphosequence development in ice-dammed (Top) and sediment-dammed basins (Bottom). The mechanism of impoundment and the chronological and topographic positions of the deposits are related to the orientation of the basins relative to the direction of ice retreat. These relationships are reflected in the organization and color coding of the List of Map Units (after Stone and others, 2005).

Postglacial Deposits (flood-plain alluvium and swamp deposits), but also including stream-terrace, talus, dune, tidal-marsh, channel fill, marine delta deposits, and artificial fill) are less widely distributed and are typically thinner than the glacial deposits that are more pervasive. The oldest postglacial deposits in Louisiana consist of coastal and poorly drained flood-plain deposits, which are key ecological elements of coastal and wetland systems. Flood-plain deposits of Tertiary alluvium are large, composed of sands, gravels and silts that have been reworked from glacial deposits and mixed with organic matter which increases their fertility. Despite their flood-

Postglacial deposits provide locally important ecological, agricultural, commercial,

QUATERNARY GEOLOGY DATA – Quaternary Geology shown on this map are from the Quaternary Geology Poly, Point Feature, and Line Feature dataset intended to be used at 1:24,000 scale. Based on Connecticut Quaternary Geology digital spatial data published in 2005 by the U.S. Geological Survey, in cooperation with the Connecticut Department of Environmental Protection. These data were digitized from the 1:24,000-scale compilation sheets prepared for the states of Connecticut. Quaternary Geology Map of Connecticut, (Stone, J.R., Schaler, J.P., London, E.H., D'Giacomo-Cohen, M. L., Lewis, R.S. and Thompson, W.B., 2005, U.S. Geological Survey special map 2 sheets, scale 1:125,000).


RELATED INFORMATION
This map is intended to be printed at its original dimensions, (48 x 36 in), in order to maintain the 1:24,000 scale (1 inch = 2,000 feet).

QUATERNARY GEOLOGY AND SURFICIAL MATERIALS DATA – 1:24,000-scale digital spatial data of Connecticut Quaternary Geology and Surficial Materials combined into one dataset, published by the Connecticut Department of Environmental Protection, in cooperation with the U.S. Geological Survey. These data were digitized from the 1:24,000-scale compilation sheets prepared for both the

BASE MAP DATA - Based on data originally from 1:24,000-scale USGS 7.5 minute topographic quadrangle maps published between 1969 and 1992. It includes political boundaries, railroads, airports, hydrography, geographic names and geographic places. Streets and street names are from Tele Atlas[®] copyrighted data. Base map information is neither current nor complete.

CONTOUR DATA - Derived from Connecticut's 2000 statewide LIDAR (Light Detection And Ranging), dataset by the University of Connecticut, College of Engineering and Natural Resources, Department of Geomatics Engineering, and the Connecticut Geological and Natural History Survey, USGS, and others. These data are a Beta product intended for research and demonstration purposes. NOTE: Contour line data is known to be incorrect in some areas due to limitations in the National Wetlands Inventory and other specific contour lines. Areas where contour lines are too straight or angular, do not naturally curve where expected, or don't exist where they probably should, do not provide indications of contour data quality.


State Plane Coordinate System of 1983, Zone 3526
Lambert Conformal Conic Projection
North American Datum of 1983



STATE OF CONNECTICUT
DEPARTMENT OF
ENVIRONMENTAL PROTECTION
79 Elm Street
Hartford, CT 06106-5127

Map created by CT DEP
December 2010

Map is not colorfast
Protect from light and moisture



WATER DEPOSITS - late Wisconsin

- Inland Melwater Deposits
- Melwater-Dammed Lakes
- Melwater-Sedimented-Dammed Lakes

Related Series of Major Ice-Dammed Ponds

- Proximal Melwater Streams
- Distal Melwater Streams

AQ DEPOSITS - late Wisconsin, Illinoian

- Deposits
- Deposits
- Deposits

EXPLANATION OF MAP SYMBOLS


- Ice Margin Position
- Inferred Ice Margin Position
- Esker
 - Glacial Stratification or Groove
 - Drumlin Axis or Ridge
- Melwater Channel
- Glacial Lake Spillway
- Inferred Glacial Spillway
- Location of Lower Till
- Two-Till Outcrop
- Deltatic Bedding Locality
- Weathered Bedrock Outcrops
- Radiocarbon-Dated Locality

lake bottom, and inland dune deposits); and Postglacial Deposits (flood-plain alluvium and swamp deposits, but also including stream-terrace, talus, dune, tidal-marsh, beach, channel fill, marine delta deposits, and artificial fill) that were emplaced in comparable topographic and depositional settings, and therefore share similar characteristics, are categorized and color coded in the Legend Description. Related Map Elements include eskers, drumlin axes, ice-margin positions, scarps, drainage divides, glacial lake spillways, meltwater channels, striations/grooves, dated sample locations, glaciofluvial and lake-bottom facies as overlays on glacial lake map units and various types of exposures.

makes them poorly drained, difficult to dig in or plow, mediocre sources of groundwater and unsuited for septic systems. Till blankets the bedrock surface in variable thicknesses and commonly underlies stratified meltwater deposits. Even moraine deposits (primarily ablation till) occur principally in northeastern Connecticut. Ice-laid deposits are inferred to be of Wisconsinan age except where exposures of older (probably Illinoian) till are shown. Drumlins are inferred to be composed of older till mantled by younger till.

Glacial Meltwater Deposits (sorted and stratified deltaic, river bottom, lake systemically, and inland pond deposits) were laid down in glacial streams, lakes and ponds which occupied the valleys and lowlands of Connecticut as the last ice sheet systemically melted away. These deposits melted away to the north. They are often composed of layers of well-sorted to poorly sorted sands, gravels, silts and clays with few or no boulders, and owing to their water-related depositional origins they have many characteristics that

The mapping presented here and on the Quaternary Geology Map of Connecticut and Long Island Sound Basin is based on recognizing single bodies of sediment or assemblages of glacial sedimentary facies that can be identified as mappable units known as morphosquences (Kotfel and Pessl, 1981). Different sedimentary facies are associated with different depositional settings and different stages of glacial retreat, and are emplaced in high-energy settings or at near the ice front. Energy levels drop off with distance from the glacier (distally) and grain size decreased along the path of meltwater flow. As a result, morphosquences are coarse grained at their collapsed, ice-contact heads and become finer distally (Figure 1). A detailed discussion of the complexities and significance of morphosquences is contained in the pamphlet that accompanies the Quaternary Geology Map of Connecticut and Long Island Sound Basin.



Forms, grading from ice-contact forms (eskers, kames) to non-ice-contact forms (flat valley graded to a specific base level. Grain-size decreases from coarse gravel at ice-contact heads, to fine sand and silt at the toe of the deposit (after Stone and others, 2005).

northward, a succession of lower bedrock spillways were opened and the valleys widened. In this case, the youngest depositional sequences occupied the lowest, widest portions of the valley (Figure 2).

and recreational resources. Talus, a result of rockfall at the base of steep bedrock (primarily trap rock) cliffs, and inland dune deposits, that developed as winds swept across newly exposed glacial lake beds, provide ecological niches that are atypical for Connecticut. Beach, dune, marsh and swamp deposits are key ecological elements of coastal and poorly drained inland settings. Deposits of floodplain alluvium are largely composed of sands, gravels and silts that have been reworked from glacial deposits and mixed with organic matter which increases their fertility. Despite their flood-prone nature, low, flat, fertile floodplains have historically been attractive for agricultural uses and development related to water-dependent commerce.

RELATED INFORMATION
This map is intended to be printed at its original dimensions, (48 x 36 in), in order to maintain the 1:24,000 scale (1 inch = 2,000 feet).

QUATERNARY GEOLOGY AND SURFICIAL MATERIALS DATA - 1:24,000-scale digital spatial data of Connecticut Quaternary Geology and Surficial Materials combined into one dataset, published by the Connecticut Department of Environmental Protection, in cooperation with the U.S. Geological Survey. These data were digitized from the 1:24,000-scale compilation sheets prepared for both the Surficial Materials Map of Connecticut, (Stone, J.R., Schaffer, J.P., London, E.H. and Thompson, W.B., 1992, U.S. Geological Survey Special Map, 2 sheets, scale 1:24,000), and map and pamphlet 71 p.) and the Connecticut Quaternary Geology Map of Connecticut, (London, E.H., Stone, J.R., Schaffer, J.P., London, E.H., DiGiacomo-Cohen, M.L., Lewis, R.L., and Thompson, W.B., 2005, U.S. Geological Survey Scientific Investigation Map 2784, 2 sheets, scale 1:25,000).

OTHER GEOLOGIC MAPS - This map is also available for individual USGS topographic quadrangles of Connecticut. This map is intended to be used with other bedrock, surficial, and quaternary (glacial) geology town maps and reports published by the Connecticut Geological and Natural History Survey, USGS, and others. Those maps and reports are also available from CT DEP.

MAPS AND DIGITAL DATA - Go to the CT ECO website for this map and a variety of others. Go to the CT DEP website for the digital spatial data shown on this map.

State Plane Coordinate System of 1983, Zone 3526
Lambert Conformal Conic Projection
North American Datum of 1983



