

QUATERNARY GEOLOGY STRATFORD, CONNECTICUT

LIST OF MAP UNITS

<p>POSTGLACIAL DEPOSITS - late Holocene, late Wisconsinan</p> <ul style="list-style-type: none"> Artificial Fill Coastal Beach and Dune Deposits Tidal Marsh Deposits Floodplain Alluvium Swamp Deposits Talus <p>EARLY POSTGLACIAL DEPOSITS - early Holocene, late Wisconsinan</p> <ul style="list-style-type: none"> Stream Terrace Deposits Inland Dune Deposits 	<p>GLACIAL MELT-WATER DEPOSITS - late Wisconsinan</p> <ul style="list-style-type: none"> Undifferentiated Meltwater Deposits Deposits of Major Ice-Dammed Lakes Deposits of Major Sediment-Dammed Ponds Deposits of Related Series of Major Ice-Dammed Ponds Deposits of Proximal Meltwater Streams Deposits of Distal Meltwater Streams <p>GLACIAL ICE-LAKE DEPOSITS - late Wisconsinan, Illinoian</p> <ul style="list-style-type: none"> Thin Till Deposits Thick Till Deposits End Moraine Deposits 	<p>Explanation of Map Symbols</p> <ul style="list-style-type: none"> Ice Margin Position Inferred Ice Margin Position Esker Glacial Striation or Groove Drumlin Axis and Center Meltwater Channel Glacial Lake Spillway Inferred Glacial Spillway Location of Lower Till Two-Till Outcrop Delatic Bedding Locality Weathered Bedrock Outcrop Radiocarbon-Dated Locality
---	---	---

EXPLANATION

Quaternary Geology is a 1:24,000-scale data that illustrates the geologic features formed in Connecticut during the Quaternary Period, which spans from 2.588 ± 0.005 million years ago to the present and includes the Pleistocene (glacial) and Holocene (postglacial) Epochs. The Quaternary Period has been a time of development of many details of the Connecticut landscape and all surficial deposits. At least twice in the last Pleistocene, continental ice sheets swept across Connecticut from the north. Their effects are of pervasive importance to present-day occupants of the land.

The Quaternary Geology information illustrates the geologic history and the distribution of depositional environments during the emplacement of unconsolidated glacial and postglacial surficial deposits and the landforms resulting from those events in Connecticut. These deposits range from a few feet to several hundred feet in thickness, overlie the bedrock surface and underlie the organic soil layer of Connecticut. Quaternary Geology is mapped without regard for any organic soil layer that may overlie the deposit.

The Connecticut Quaternary Geology information was initially compiled at 1:24,000 scale (1 inch = 2,000 feet) for incorporation for a statewide 1:125,000-scale map, Quaternary Geology Map of Connecticut and Long Island Sound Basin. A companion map, the Surficial Materials Map of Connecticut, emphasizes the surface and subsurface texture (grain-size distribution) of these materials. The quaternary geology and surficial material features portrayed on these two maps are very closely related, each contributing to the interpretation of the other.

Most of Connecticut's surficial material is glacially derived, and can be divided into two broad depositional categories: Glacial Ice-Lake Deposits (nonsorted and generally nonstratified till, thick till, and end moraine) which are generally exposed in the uplands, and are the most widespread surficial deposit in Connecticut; and Glacial Meltwater Deposits (sorted and stratified delatic, river bottom, lake bottom, and inland dune deposits) which are most commonly concentrated in valleys and lowlands.

Particular attention has been paid to understanding the distribution and characteristics of stratified meltwater deposits because they have historically influenced development patterns and groundwater availability throughout the state. Within the meltwater category, six classes of deposits have been recognized based on the conditions that prevailed during their emplacement: various depositional settings, including previously deposited sediment, or the glacier itself, impounded the lake or pond where emplacement occurred (see the meltwater deposit discussion below). Meltwater stream deposits are differentiated based on their distance (proximal or distal) from the ice sheet when they were emplaced, and a separate meltwater map unit is reserved for deposits of undetermined provenance (unaccounted).

Postglacial Deposits were emplaced by various processes after the melt back of the last ice sheet. Some of these deposits were emplaced early in post-glacial time and have been grouped together as Early Postglacial Deposits. Later deposits, resulting from processes that are still active (or are manmade), have been grouped together as Postglacial Deposits.

Glacial Ice-Lake Deposits (nonsorted and generally nonstratified till, thick till, and end moraine), Glacial Meltwater Deposits (sorted and stratified delatic, river bottom, lake bottom, and inland dune deposits), and Glacial Meltwater Deposits (sorted and stratified delatic, river bottom, lake bottom, and inland dune deposits) are generally exposed in the uplands, and are the most widespread surficial deposit in Connecticut, and Glacial Meltwater Deposits (sorted and stratified delatic, river bottom, lake bottom, and inland dune deposits) which are most commonly concentrated in valleys and lowlands.




Figure 1: A morphosequence is a body of meltwater deposits composed of a continuum of land forms, grading from ice-contact forms (eskers, kames) to non-ice-contact forms (flat valley terrace, delta plains), that were deposited simultaneously at and beyond the margin of a glacier, graded to a specific base level. Grain-size decreases faster course grain at ice-contact, roads, through sand and gravel and sand beneath delta plains and forest slopes to silt and clay in lake-bottom deposits (after Stone and others, 2005).

Deposition of the morphosequences that progressively filled bedrock valleys and lowlands as the last glacier melted northward required the presence of impounded lakes and ponds. The nature of the impoundment, and the oldest morphosequence were emplaced in the higher, narrower portions of the basin. As the ice front retreated northward, a succession of lower bedrock spillways were opened and the valleys widened. In this case, the youngest depositional sequences occupied the lowest, widest portions of the valley (Figure 2).

DATA SOURCES

QUATERNARY GEOLOGY DATA - Quaternary Geology shown on this map are from the Quaternary Geology Poly, Point Feature, and Line Feature dataset intended to be used at 1:24,000 scale. Based on Connecticut Quaternary Geology digital spatial data published in 2005 by the U.S. Geological Survey, in cooperation with the Connecticut Department of Environmental Protection. These data were digitized from the 1:24,000-scale compilation sheets prepared for the statewide Quaternary Geology Map of Connecticut, (Stone, J.R., Schaller, J.P., London, L.H., DiGiacomo-Cohen, M.L., Lewis, R.S., and Thompson, W.B., 2005, U.S. Geological Survey special map, 2 sheets, scale 1:125,000).

BASE MAP DATA - Based on data originally from 1:24,000-scale USGS 7.5 minute topographic quadrangle maps published between 1969 and 1992. It includes political boundaries, railroads, airports, hydrography, geographic names and geographic places. Streets and street names are from Tele Atlas' copyrighted data. Base map information is neither current nor complete.

CONTOUR DATA - Derived from Connecticut's 2000 statewide LIDAR (Light Detection And Ranging) dataset by the University of Connecticut, College of Agriculture and Natural Resources, Department of Natural Resources and the Environment. These data are a beta product intended for research and demonstration purposes. NOTE: Contour line data is known to be incorrect in some areas due to anomalies in the underlying elevation data used to generate those specific contour lines. Areas where contour lines are too straight or angular, do not naturally curve where expected, or don't exist where they probably should are good indications of erroneous data.

