

ICE-LAID DEPOSITS

- Thin Till
- Thick Till
- End Moraine deposits

ACIAL MELT-WATER DEPOSITS

ane Deposits

- Fines (very fine sand, silt, and clay)

arse Deposits

- Gravel
- Sand and Gravel
- Sand

acked Coarse Deposits

- Gravel overlying Sand and Gravel
- Gravel overlying Sand
- Sand and Gravel overlying Sand
- Sand overlying Gravel
- Sand overlying Sand and Gravel

acked Coarse Deposits Overlying Fine Deposits

- Gravel overlying Sand overlying Fines
- Gravel overlying Fines
- Sand and Gravel overlying Sand overlying Fines
- Sand and Gravel overlying Fines

acked Fine Deposits Overlying Coarse Deposits

- Fines overlying Sand and Gravel
- Fines overlying Sand

POSTGLACIAL DEPOSITS

- Floodplain Alluvium
- Alluvium overlying undifferentiated Coarse deposits (g. sq. ss)
- Alluvium overlying Sand
- Alluvium overlying Fines
- Alluvium overlying undifferentiated Coarse deposits overlying Fine deposits
- Alluvium overlying undifferentiated Fine deposits overlying Coarse deposits
- Swamp deposits
- Swamp deposits overlying Sand
- Swamp deposits overlying Fines
- Swamp deposits overlying Sand overlying Fines
- Swamp deposits overlying Fines overlying Sand
- Salt-Marsh and Tidal-Marsh deposits
- Salt-Marsh and Tidal-Marsh deposits overlying Sand
- Salt-Marsh and Tidal-Marsh deposits overlying Fines
- Salt-Marsh and Tidal-Marsh deposits overlying Fines
- Talus
- Beach deposits
- Artificial Fill

* Alluvium may be overlying any of the Coarse deposits (g. sq. ss)

w Water

		PARTICLE DIAMETER															
		10	2.5	.16	.08	.04	.02	.01	.005	.0025	.0015	in					
		256	64	2	1	5	25	125	625	2048		mm					
Boulders	Coarse	Pebbles	Granules	Sand	Very Coarse Sand	Medium Sand	Fine Sand	Very Fine Sand	Silt	Clay	City						
	GRAVEL PARTICLES				SAND PARTICLES				FINE PARTICLES								

Grain-size classification (modified from Wentworth, 1922)

Unconsolidated glacial and postglacial deposits, that range from a few feet to several hundred feet in thickness, overlie the bedrock surface of Connecticut (see Block Diagram). This map portrays the distribution of these deposits. The legend is designed to highlight the relationship between the depositional origins and the composition and character of the materials portrayed. Most of the Connecticut glacial and postglacial deposits can be divided into two broad depositional categories: Glacial Ice-Land deposits (tills and moraine) which are generally deposited on the land surface by glacial meltwater, and Glacial Water-Land deposits (stratified deposits) which are most commonly concentrated in valleys and lowlands. A mapping emphasis is placed on stratified meltwater deposits, which are the most important and variable, and historically influenced development patterns throughout the state.

Glacial Ice-Laid Deposits (tills and moraine) were derived directly from the ice and consist of nonsorted, generally nonstratified mixtures of grain-sizes ranging from clay to large boulders. The tills and moraine deposits are predominantly composed of sand and silt that can be sparse to abundant. Some tills contain lenses of sorted sand and gravel and occasionally masses of laminated fine-grained sediment. The lack of sorting and stratification typical of ice-laid deposits is predominantly a function of the rapid deposition of meltwater, mediceous sources of groundwater and unsieved for septic systems. Till blankets the bedrock surface in variable thicknesses and commonly underlies stratified meltwater deposits (see Block 10). The primary alluvial till is primarily alluvial till occurs principally in southeastern Connecticut.

SURFICIAL MATERIALS DATA – Surficial Materials shown on this map are from the Surficial Material Poly dataset which contains polygon data intended to be used at a 1:24,000 scale. Based on Connecticut Department of Environmental Protection, 1998, *Surficial Materials Digital Data*, prepared by Connecticut Department of Environmental Protection, in cooperation with the U.S. Geological Survey. These data were digitized from the 1:24,000-scale compilation sheets prepared for the statewide Surficial Materials Map of Connecticut. (Stone, J.R., 1998, *Surficial Materials Map of Connecticut*, U.S. Geological Survey special map, 2 sheets, scale 1:125,000).

QUATERNARY GEOLOGY AND SURFICIAL MATERIALS DATA - 1:24,000-scale digital spatial data of Connecticut Quaternary Geology and Surficial Materials combined into one dataset. Data were compiled by Connecticut Department of Environmental Protection, in cooperation with the U.S. Geological Survey. These data were digitized from the 1:24,000-scale compilation sheets prepared for both the Surficial Materials Map of Connecticut, Stone and others, 1998, 1:125,000 and the Quaternary Geology Map of Connecticut and Long Island Sound Basin, Stone and others, 2005, 1:125,000.

RELATED INFORMATION
This map is intended to be printed at its original dimensions in order to maintain the 1:24,000 scale (1 inch = 2000 feet).

MAP LOCATION

1 0.5 0 1 Miles

1000 500 0 500 1000 Feet

1 Kilometers

SCALE 1:24,000 (1 inch = 2000 feet) when map is printed at original size

Melwater deposits are depicted using four basic texturally-based melt unit types: gravel, sand and gravel, sand, and fines. To the extent that it is known or can be inferred, the subsurface textural composition of meltwater deposits is shown for their entire vertical extent. The time that a meltwater deposit was being laid down, and a single map unit (e.g. s-sd) is sufficient to describe the entire meltwater section. Areal and vertical textural variability can occur within the meltwater section, but is not shown here. Energy away from the sedimentation varies with each meltwater setting (melt, delta, lake, etc.), and settings can change over time. High-energy depositional environments near glacial margins (proximal) tend to favor deposition of coarse material but, as time passes, and the glacial margin retreats, meltwater settings change and the deposition of finer deposits can become predominant. Where more complex stratigraphic relationships existed because of changing conditions during deposition, "stacked" map units are used to characterize the subsurface (e.g. sg-sd and gravel overlying sand overlying fines). The relationship is also shown (e.g. g-sd - alluvium overlying sand).

SOURCES

QUATERNARY GEOLOGY AND SURFICIAL MATERIALS. DATA - 1:24,000-scale digital spatial data of Connecticut Quaternary Geology and Surficial Materials combined into one dataset, published by the Connecticut Department of Environmental Protection, in cooperation with the U.S. Geological Survey. These data were digitized from the 1:24,000-scale compilation sheets prepared for both the Surficial Materials Map of Connecticut, Stone and others, 1992, 1:125,000 and the Quaternary Geologic Map of Connecticut and Long Island Sound Basin, Stone and others, 2005, 1:125,000.

MAPS AND DIGITAL DATA - Go to the CT ECO website for this map and a variety of others. Go to the CT DEP website for the digital spatial data shown on this map.

A map of Ohio showing county boundaries. A black rectangle in the central part of the state indicates the study area. To the left of the map is a scale bar labeled '1 Miles'. To the right of the map is a north arrow pointing upwards.

State Plane Coordinate System of 1983, Zone 35
Lambert Conformal Conic Projection
North American Datum of 1983

by CT DEP
2009
colorfast
and moisture

 **USGS**
science for a changing world

