

**ACIAL ICE-LAID DEPOSITS**

- Thin Till
- Thick Till
- End Moraine deposits

**ACIAL MELT-WATER DEPOSITS**

**Desposits**

- Fines (very fine sand, silt, and clay)

**Coarse Deposits**

- Gravel
- Sand and Gravel
- Sand

**acked Coarse Deposits**

- Gravel overlying Sand and Gravel
- Gravel overlying Sand
- Sand and Gravel overlying Sand
- Sand and Gravel overlying Sand overlying Sand and Gravel
- Sand overlying Gravel
- Sand overlying Sand and Gravel





**acked Coarse Deposits Overlying Fine Deposits**

- Gravel overlying Sand overlying Fines

**POSTGLACIAL DEPOSITS**

- Floodplain Alluvium
- Alluvium overlying undifferentiated Coarse deposits (g, s, s)
- Alluvium overlying Sand
- Alluvium overlying Fines
- Alluvium overlying undifferentiated Coarse deposits overlying Fine deposits
- Alluvium overlying undifferentiated Fine deposits overlying Coarse deposits
- Swamp deposits
- Swamp deposits overlying Sand
- Swamp deposits overlying Fines
- Swamp deposits overlying Sand overlying Fines
- Swamp deposits overlying Fines overlying Sand
- Salt-Marsh and Tidal-Marsh deposits
- Salt-Marsh and Tidal-Marsh deposits overlying Sand
- Salt-Marsh and Tidal-Marsh deposits overlying Fines
- Talus
- Beach deposits
- Artificial Fill

\* Alluvium may be overlying any of the Coarse deposits (g, s, s)


	Sand and Gravel overlying Sand overlying Fines
	Sand and Gravel overlying Fines
	Sand overlying Fines
<b>Packed Fine Deposits Overlying Coarse Deposits</b>	
	Fines overlying Sand and Gravel
	Fines overlying Sand

Unconsolidated glacial and postglacial deposits, that range from a few feet to several hundred feet in thickness, overlie the bedrock (see Fig. 1). These deposits are composed of a wide range of the areal extent, subsurface, grain-size, and textural distributions of these surficial materials. The current research is designed to highlight the relationship between the depositional origins and the distribution of the character of the materials portrayed. Most of Connecticut's surficial material is glacially derived, and can be divided into two broad depositional categories: Glacial Ice-land deposits (fills and moraine) which are generally exposed in the western half of the state, and Glacial Melwater deposits (stratified deposits), which are most commonly concentrated in valleys and lowlands. A major ongoing research project is to determine the depositional influence their distribution and character have historically influenced depositional patterns throughout the state.

For a complete description of surficial materials map units, and further information concerning their thickness and modes of occurrence, please refer to the published Surficial Materials Map of Connecticut and the companion Quaternary Geologic Map of Connecticut and Long Island Sound Basin (see Data Sources).


**Glacial Ice-Laid deposits** (tills and moraine) were derived directly from the ice and consist of nonsorted, generally nonstratified mixtures of grain-sizes ranging from clay to large boulders. The matrix of most tills is predominantly sand and silt and boulders can be sparse to abundant. Some tills contain lenses of sorted sand and gravel and occasionally masses of laminated fine-grained sediment. The lack of sorting and stratification typical of ice-laid deposits often makes them poorly drained, difficult to dig in, and places a medium-sized boulder under and unsuited for septic systems. Till blankets the bedrock surface in variable thicknesses and commonly underlies stratified meltwater deposits (see Block Diagram). End moraine deposits (primarily albion till) occur principally in southeastern Connecticut.

**Glacial Meltwater deposits** (stratified deposits) were laid down in glacial streams, lakes and ponds which occupied the valleys and lowlands of Connecticut as the last ice sheet melted away to the north. They are often composed of layers of well-to-poorly sorted sands, gravels, silts and clays with few to no boulders, and owing to their water-related depositional origins they have many



EXPLANATION OF CROSS-SECTION VIEW

EXPLANATION OF SURFACE VIEW




**SURFICIAL MATERIALS DATA** – Surficial Materials shown on this map are from the Surficial Material Poly dataset which contains polygon data intended to be used at 1:24,000 scale. Based on Connecticut Surficial Materials digital data published in 1995 by the Connecticut Department of Environmental Protection, in cooperation with the U.S. Geological Survey. These data were digitized from the 1:24,000 scale compilation sheets prepared for the statewide Surficial Materials Map of Connecticut, (Stone, J.R., Schafer, J.P., London, E.H. and Thompson, W.B., 1992, U.S. Geological Survey special map, 2 sheets, scale 1:125,000).

**BASE MAP DATA** - Based on data originally from 1:24,000-scale USGS 7.5 minute topographic quadrangle maps published between 1969 and 1992. It includes political boundaries, railroads, airports, hydrography, geographic names and geographic places. Streets and street names are from Tele Atlas® copyrighted data. Base map information is neither current nor complete.

### RELATED INFORMATION

This map is intended to be printed at its original dimensions in order to maintain the 1:24,000 scale (1 inch = 2000 feet).



characteristics that are favorable for development. Because water is a better sorting agent than ice, glacial meltwater deposits are commonly better sorted, more permeable, and better aquifers than ice-laid deposits. They can be good sources of construction aggregate, and are relatively easy to excavate and build highways and buildings on.

Meltwater deposits are depicted using four basic texturally-based map units: gravel, sand and gravel, sand, and fines. To the extent that it is known or can be inferred, the subsurface textural characteristics of these units are indicated by the thickness of the line representing the unit. In many places similar conditions persisted for the entire time that a meltwater deposit was being laid down, and a single map unit (e.g. sand) is sufficient to describe the entire meltwater sequence. Areal and vertical textural variability can occur within the meltwater sequence, and this is indicated by the use of the same map unit in different sediment varies with each meltwater setting (stream, delta, lake, etc.), and settings can change over time. High-energy depositional environments near glacial margins (proximal) tend to favor coarser-grained deposits, and these deposits are more likely to be preserved. In more distal settings, meltwater beds, less energy is available and finer grained silted deposits can become predominant. Where more complex stratigraphic relationships existed because of changing conditions during deposition, "stacked" map units are used to characterize the meltwater sequence. The use of the same map unit for different meltwater lines. Where postglacial deposits overlie meltwater deposits, this relationship is also shown (e.g. as -/ overlying sand).

**Postglacial Sediments** (primarily floodplain alluvium and swamp deposits) are less widely distributed and are typically thinner than the glacial deposits that they overlie, but they are locally important ecological, agricultural, commercial, and recreational resources. Talus, a result of rockfall at the base of steep bedrock (primarily trap rock) cliffs, provides a cool damp ecological niche. Beach, marsh and swamp deposits are key ecological elements of coastal and poorly drained inland settings. Deposits of floodplain alluvium

are largely composed of sands, gravels and silts that have been reworked from glacial deposits and mixed with organic matter which increases their fertility. Despite their flood-prone nature, low, flat, fertile floodplains have historically been attractive for agricultural uses and development related to water-dependent commerce.

This geological cross-section illustrates the subsurface structure of the study area. It shows a sequence of geological units, including the Quaternary (Q), Tertiary (T), and Cretaceous (C) formations. The units are color-coded and labeled with their respective names and codes. The cross-section also shows the topography of the area, with a prominent mountain range in the center. The units are shown in a stratigraphic column on the right, with the Quaternary at the top, followed by the Tertiary, and the Cretaceous at the bottom. The units are labeled with their respective names and codes: Q (Quaternary), T (Tertiary), and C (Cretaceous). The units are shown in a stratigraphic column on the right, with the Quaternary at the top, followed by the Tertiary, and the Cretaceous at the bottom. The units are labeled with their respective names and codes: Q (Quaternary), T (Tertiary), and C (Cretaceous).

EXPLANATION OF SURFACE VIEW

travel deposits      s/f      Sand overlying fines

is	f/g	Fines overlying sand and gravel
lying sand	s	Alluvium
avel overlying sand		Alluvium overlying fines
ing sand and gravel	t	Till and bedrock

gravel overlying sand overlying fines       Thick till

gravel overlying fines       Thin till

## SOURCES

Quaternary Geology and Surficial Materials DATA - 1:24,000-scale digital spatial data of Connecticut Quaternary Geology and Surficial Materials combined into one dataset, published by the Connecticut Department of Environmental Protection, in cooperation with the U.S. Geological Survey. These data were digitized from the 1:24,000-scale

**OTHER GEOLOGIC MAPS** - This map is also available for purchase from the U.S. Geological Survey, 600 G Street, NW, Washington, DC 20004, 1-800-354-5434.

bedrock, surficial, and quaternary (glacial) geology quadrangle maps and reports published by the Connecticut Geological and Natural History Survey, USGS, and others are also available from CT DEP.


**MAPS AND DIGITAL DATA** - Go to the CT ECO website for

MAP LOCATION

State Plane Coordinate System of 1983, Zone 35  
Lambert Conformal Conic Projection  
North American Datum of 1983

USGS  
science for a changing world

---



STATE OF CONNECTICUT  
DEPARTMENT OF  
ENVIRONMENTAL PROTECTION  
79 Elm Street  
Hartford, CT 06106-5127

Map created by CT DEP  
August 2009  
Map is not colorfast  
Protect from light and moisture

