

ICE-LAKE ICED LAID DEPOSITS

- Thin Till
- Thick Till
- End Moraine deposits

GLACIAL MELT-WATER DEPOSITS

Clay Deposits

- Fines (very fine sand, silt, and clay)

Sand Deposits

- Gravel
- Sand and Gravel
- Sand

Unsorted Coarse Deposits

- Gravel overlying Sand and Gravel
- Gravel overlying Sand
- Sand and Gravel overlying Sand
- Sand and Gravel overlying Sand overlying Sand and Gravel
- Sand overlying Gravel
- Sand overlying Sand and Gravel

Sorted Coarse Deposits Overlying Fine Deposits

- Gravel overlying Sand overlying Fines
- Gravel overlying Fines
- Sand and Gravel overlying Sand overlying Fines
- Sand and Gravel overlying Fines
- Sand overlying Fines

Sorted Fine Deposits Overlying Coarse Deposits

- Fines overlying Sand and Gravel
- Fines overlying Sand

POSTGLACIAL DEPOSITS

- Floodplain Alluvium
- Alluvium overlying undifferentiated Coarse deposits (s.g.s.)
- Alluvium overlying Sand
- Alluvium overlying Fines
- Alluvium overlying undifferentiated Coarse deposits overlying Fine deposits
- Alluvium overlying undifferentiated Fine deposits overlying Coarse deposits
- Swamp deposits
- Swamp deposits overlying Sand
- Swamp deposits overlying Sand
- Swamp deposits overlying Sand overlying Fines
- Swamp deposits overlying Fines overlying Sand
- Salt-Marsh and Tidal-Marsh deposits
- Salt-Marsh and Tidal-Marsh deposits overlying Sand
- Salt-Marsh and Tidal-Marsh deposits overlying Fines
- Talus
- Beach deposits
- Artificial Fill

* Alluvium may be any of the Coarse deposits (s.g.s.)

w Water

		PARTICLE DIAMETER											
		10	2.5	.16	.08	.04	.02	.01	.005	.0025	.0015	.001	mm
		256	64	4	2	1	0	0	25	125	625	3125	15625
Boundaries	Coarses	Pebbles	Cobbles	Gravel		Coarse Sand	Medium Sand	Fine Sand	Very Fine Sand		Silt	Clay	
		GRAVEL PARTICLES						SAND PARTICLES			FINE PARTICLES		

Grain-size classification (modified from Wentworth, 1922)


Unconsolidated glacial and postglacial deposits, that range from a few feet to several hundred feet in thickness, overlie the bedrock (see Fig. 1). These deposits are composed of a wide range of the areal extent, subsurface, grain-size, and textural distributions of these surficial materials. The next chapter is designed to highlight the relationship between the depositional origins and the distribution of the character of the materials portrayed. Most of Connecticut's surficial material is glacially derived, and can be divided into two broad depositional categories: Glacial Ice-land deposits (fills and moraine) which are generally deposited in the central and western portions of the state, and the Glacial Marine; Connecticut, and Glacial Meltdown deposits (stratified deposits) which are most commonly concentrated in valleys and lowlands. A brief description of the origin and distribution of these deposits followed by their distribution and character have historically influenced depositional patterns throughout the state.

Glacial Ice-Laid Deposits (tilts and moraine) were derived directly during the ice contact and consist of nonsorted, generally nonstratified, unconsolidated, noncemented, and nonlaminated material. The composition of most tills is predominantly sand and silt and boulders can be locally abundant. The moraine deposits are composed of sand and gravel and occasionally masses of laminated fine-grained sediment. The lack of sorting and stratification typical of ice-laid deposits is a result of the rapid deposition of material from a meltwater flow, meltwater sources of groundwater and unsuitable for septic use. The moraine deposits are typically composed of sand and gravel and commonly underlie stratified meltwater deposits (see Block Diagram). End moraine deposits (primarily ablation till) occur in the northern part of the study area.

During deposition, "stacked" map units are used to characterize the subsurface (e.g. gs/s) sand and gravel overlying sand overlying silt and clay. The relationship is also shown (e.g. a/s - alluvium overlying sand).

Postglacial Sediments (primarily floodplain alluvium and swamp deposits) are less widely distributed and are typically thinner than the glacial deposits. They are composed of sand, silt, and clay, and are used for agricultural, commercial, and recreational purposes. The deposits are typically composed of sand and gravel and commonly underlie stratified meltwater deposits (see Block Diagram). End moraine deposits (primarily ablation till) occur in the northern part of the study area.

Melwater deposits are depicted using four basic texturally-based map units: gravel, sand and gravel, sand, and fines. To the extent that it is known or can be inferred, the subsurface texture and thickness of these units is indicated by the use of different line thickness. In many places similar conditions persisted for the entire time that a melwater deposit was being laid down, and a single map unit (e.g. s-sand) is sufficient to describe the entire melwater sequence. Areal and vertical texture/variability can occur within the melwater sequence, and this is indicated by the use of different line patterns. Sediment varies with each melwater setting (stream, delta, lake, etc.), and settings can change over time. High-energy depositional environments near glacial margins (proximal) tend to favor coarser textures, and as the distance from the glacial margin increases, less energy is available and finer grained silt/clay textures become predominant. Where more complex stratigraphic relationships existed because of changing conditions during deposition, "stacked" map units are used to characterize the melwater sequence. The use of the letters "a" and "b" in the map units (e.g. a-s-s) where postglacial deposits overlie melwater deposits, this relationship is also shown (e.g. a/s - alluvium overlying sand).



SURFICIAL MATERIALS DATA – Surficial Materials shown on this map are from the Surficial Material Poly dataset which contains polygon data intended to be used at 1:24,000 scale. Based on the Connecticut State Survey of Geologic Data, compiled by the Connecticut Department of Environmental Protection, in cooperation with the U.S. Geological Survey. These data were digitized from the 1:24,000-scale compilation sheets prepared for the Connecticut State Survey of Geologic Data, 1959-1965, by J.P. Schafer, J.P., London, E.H. and Thompson, W.B., 1992, U.S. Geological Survey special map, 2 sheets, scale 1:125,000.

RELATED INFORMATION
This map is intended to be printed at its original dimensions in order to maintain the 1:24,000 scale (1 inch = 2000 feet).

MAPS AND DIGITAL DATA - Go to the CT ECO website for this map and a variety of others. Go to the CT DEP website for the digital spatial data shown on this map.

